arXiv:1210.7163v1 [math.DS] 26 Oct 2012

INFINITESIMAL DEFORMATIONS OF RATIONAL SURFACE AUTOMORPHISMS

by

Julie Déserti & Julien Grivaux

Abstract. — This article deals with families of automorphisms of rational surfaces and related rigidity properties. We introduce the notions of rigidity and infinitesimal rigidity, and give a bound on the number of parameters of a family of rational surfaces endowed with an automorphism. This yields in particular a cohomological criterion to prove the rigidity of an automorphism. We study automorphisms on rational surfaces X whose anticanonical divisor is effective but also automorphisms of rational Kummer surfaces. Finally we construct a theoretical framework to compute effectively in any concrete example the action of a rational surface automorphism on the space of infinitesimal deformations of the surface.

2010 Mathematics Subject Classification. — 37F10, 14E07, 32G05

Contents

1. Introduction 2
2. Infinitesimal rigidity 3
    2.1. Infinitesimal rigidity is a spectral property 3
    2.2. Automorphisms of anticanonical basic rational surfaces 4
3. A geometric example: Kummer surfaces 6
    3.1. Construction, Picard group and fixed points 6
    3.2. An alternative construction via foliations 10
    3.3. Action of the automorphism group on infinitesimal deformations 10
4. Realization of infinitesimal deformations using divisors 12
    4.1. 1-exceptional divisors 13
    4.2. Geometric bases 15
    4.3. Algebraic bases 16
5. An explicit example on $\mathbb{P}^2$ blown up in 15 points 17
    5.1. The strategy 17
    5.2. Calculations for the pair (D_1, D_2) 18
    5.3. Calculations for the pair $(\mathfrak{D}_1, \mathfrak{D}_2)$ 22
    5.4. The result 24
References 26
6. Maple computations 26

First author supported by the Swiss National Science Foundation grant no PP00P2_128422 /1.
Both authors supported by ANR Grant "BirPol" ANR-11-JS01-004-01.

1. Introduction

Biregular automorphisms of rational surfaces with positive topological entropy present a major interest in complex dynamics, but their construction remains a difficult problem of algebraic geometry. For an overview of this problem, we refer to [**7**] and to the references therein. It is a paradoxal fact that these automorphisms, although hard to construct, can occur in holomorphic families of arbitrary large dimension (*see* [**2**]). Besides, the automorphism group of a given rational surface can carry many automorphisms of positive entropy; we refer to [**4**] and [**3**] for recent and interesting results on this topic. In this paper, pursuing our previous work [**7**], we concern ourselves with families of automorphisms of rational surfaces and related rigidity properties.

Let $\mathfrak{X} = (X_t)_{t \in B}$ be a holomorphic family of complex manifolds parameterized by a smooth base B. The theory of deformations allows us to associate to $\mathfrak{X}$ a number $\mathfrak{m}(\mathfrak{X})$ of generic parameters between 0 and $\dim B$ that measures the variation of the complex structure of the generic fiber of the family $\mathfrak{X}$ (*see* [**7**]). We say that a biholomorphism f of a complex manifold X is *rigid* if any deformation $\mathfrak{X}$ of X on which the automorphism f can be extended satisfies $\mathfrak{m}(\mathfrak{X}) = 0$. If f is an automorphism of X that can be extended to a deformation $\mathfrak{X}$, then the Kodaira-Spencer map at the central fiber commutes to the action f^* of f on $\mathrm{H}^1\big(X, \mathrm{T}X\big)$; therefore the image of $\mathrm{KS}(\mathfrak{X})$ is fixed by this action. In particular $\mathfrak{m}(\mathfrak{X}) \leq \dim\ker(f^* - \mathrm{id})$; this inequality allows to give on the one hand a upper bound for $\mathfrak{m}(\mathfrak{X})$ and on the other hand a simple criterion to say whether (X, f) is rigid:

Theorem A. — *Let f be an automorphism of X and let f^* be the corresponding action of f on $\mathrm{H}^1\big(X, \mathrm{T}X\big)$. If $\mathfrak{X}$ is a deformation of X such that f lifts to an automorphism of $\mathfrak{X}$, then*

$$\mathfrak{m}(\mathfrak{X}) \leq \dim\ker(f^* - \mathrm{Id}).$$

In particular, if 1 is not an eigenvalue of f^* then f is rigid.

An automorphism f of X is *infinitesimally rigid* if f cannot be lifted to an automorphism of a non-trivial infinitesimal deformation of X, or equivalently if 1 is not an eigenvalue of f^* acting on $\mathrm{H}^1\big(X, \mathrm{T}X\big)$. Remark that an automorphism can be rigid without being infinitesimally rigid, but the converse holds. The main problem in order to apply Theorem A in practical examples is that the computation of the action of $\mathrm{Aut}(X)$ on $\mathrm{H}^1\big(X, \mathrm{T}X\big)$, this is more delicate than the action of $\mathrm{Aut}(X)$ on the Picard group of X. The reason for this is that the first action is not defined for birational automorphisms, whereas the second is. Our purpose in this article is to compute this action in various examples.

The first case we deal with is the case of rational surfaces carrying an anticanonical curve. This class of rational surfaces is particularly relevant for us because automorphisms on these surfaces are well understood and studied (*see* for instance [**11**] and [**8**]). The most significant result we prove in this setting is the following:

Theorem B. — *Let X be a basic rational surface with $\mathrm{K}_X^2 < 0$ endowed with an automorphism f, and assume that there exists an irreducible singular curve C on X which is linearly equivalent to $-\mathrm{K}_X$. Then f is not infinitesimally rigid. Besides, X has no nonzero holomorphic vector field.*

Unfortunately, Theorem B doesn't allow to decide if f is rigid or not, all it says is that the rigidity of f cannot be proved using an approximation at the first order.

The second case we deal with is the case of rational Kummer surfaces. Let $\mathcal{E}$ be the elliptic curve obtained by taking the quotient of the complex line $\mathbb{C}$ by the lattice $\Lambda = \mathbb{Z}[\mathbf{j}]$ $\big($resp. $\Lambda = \mathbb{Z}[\mathbf{i}]\big)$. The group $\mathrm{GL}(2;\Lambda)$ acts linearly on the complex plane and preserves the lattice $\Lambda \times \Lambda$; therefore any element M of $\mathrm{GL}(2;\Lambda)$ induces an automorphism f_M on $\mathcal{E} \times \mathcal{E}$ that commutes with the automorphism ι (resp. ϕ) defined by $\iota(x,y) = (\mathbf{i}x, \mathbf{i}y)$ $\big($resp. $\phi(x,y) = (\mathbf{j}x, \mathbf{j}y)\big)$. The automorphism f_M lifts to an automorphism $\widetilde{f_M}$ on the desingularization X of $(\mathcal{E} \times \mathcal{E})/\langle\iota\rangle$, which is a rational Kummer surface. Besides the topological entropy of $\widetilde{f_M}$ is twice the logarithm of the spectral radius of M.

Theorem C. — *For any matrix M of infinite order in $\mathrm{GL}(2;\mathbb{Z}[\mathbf{j}])$, the automorphism $\widetilde{f_M}$ is rigid.*

The situation is drastically different for the lattice $\mathbb{Z}[\mathbf{i}]$. Indeed, we have the following statement:

Theorem D. — *For any matrix M in $\mathrm{GL}(2;\mathbb{Z}[\mathbf{i}])$ such that $(\det M)^2 = 1$, the automorphism $\widetilde{f_M}$ is not infinitesimally rigid. However, there exist elements in $\mathrm{GL}(2;\mathbb{Z}[\mathbf{i}])$ whose associated automorphism is infinitesimally rigid.*

Note that the automorphisms of rational Kummer surfaces appear in different important problems. Let us for instance recall the following result [**6**]: let f be a birational map of a projective surface that preserves a holomorphic foliation $\mathcal{F}$. If $(\deg f^n)_n$ grows exponentially, then up to birational conjugacy and to a finite covering, either f is a Anosov automorphism on $\mathbb{C}^2/(\Lambda\times\Lambda)$ with $\Lambda=\mathbb{Z}[\mathbf{j}]$ or $\Lambda=\mathbb{Z}[\mathbf{i}]$, or f is a monomial map on $\mathbb{P}^1\times\mathbb{P}^1$. Moreover $\mathcal{F}$ is linear and f preserves at least two foliations. As we will see in §3.2 these examples also appear in the so-called Poincaré Problem.

Lastly, if f is an automorphism of a basic rational surface X, we explain how to compute in practice the action of f on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ from the data of the sequence of blowups of $\mathbb{P}^2$ defining X and the explicit formula of f considered as a birational map of $\mathbb{P}^2$. For doing this, we develop specific theoretical tools, such as 1-exceptional divisors. Then we carry out completely the computation in one specific example, which is the one constructed in [**7**, Thm 3.5]; it is an automorphism obtained by blowing up 15 successive points of $\mathbb{P}^2$, which correspond to three infinitely near points of length five. Some of the computations are computer-assisted (with Maple), we provide as an appendix of this paper a commented Maple file.

Let us describe in more details the organization of the paper. In §2, we deal with infinitesimal rigidity of rational surface automorphisms. We introduce in §2.1 the notions of rigidity and infinitesimal rigidity, and we give in Theorem 2.2 a bound on the number of parameters of a family of rational surfaces endowed with an automorphism. This gives in particular a cohomological criterion to prove the rigidity of an automorphism. In §2.2, we study automorphisms on rational surfaces X whose anticanonical divisor is effective, and we compute in Theorems 2.4 and 2.5 the action of the automorphism group on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ in specific situations. In particular, we prove that a rational surface automorphism is never infinitesimally rigid if there exists a singular irreducible anticanonical curve on the surface. In §3, we deal with automorphisms of rational Kummer surfaces. Section §3.1 consists in recollections about the construction of these surfaces, as well as formulæ for the action of the automorphisms on the Picard group. We limit ourselves to the case of hexagonal lattices. Finally, using the Atiyah-Bott fixed point formula, we compute in Corollary 3.5 the character of the representation of $\mathrm{Aut}(X)$ in $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ when X is a Kummer surface of hexagonal type. In §3.2, following [**10**] and [**15**], we present another approach of the construction of rational Kummer surfaces via foliations. This approach is geometric in nature and allows to prove in an elegant way that these surfaces are indeed rational. We have tried to make a self-contained exposition of the theory for the reader's convenience, but we claim no originality in these two sections. In §3.3, we compute explicitly the characteristic polynomial of the action of an automorphism of a Kummer surface on its infinitesimal deformation space: for hexagonal lattices, this is Theorem 3.7 which refines Corollary 3.5; for cubic lattices, this is Theorem 3.9. The purpose of §4 is the construction of a theoretical framework to compute effectively in any concrete example the action of a rational surface automorphism on the space of infinitesimal deformations of the surface. The main idea is to represent the space $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ of infinitesimal deformations of X by global sections of the sheaf $\frac{\mathrm{T}X(D)}{\mathrm{T}X}$ for some special divisors D called 1-exceptional divisors, they are studied in §4.1. Then we construct two different bases of $\frac{\mathrm{T}X(D)}{\mathrm{T}X}$ in §4.2 and §4.3, both will be useful for practical computations of the action of automorphisms. The last section is devoted to an explicit example, and relies on the machinery developed in §4. The strategy is explained in §5.1, and the computations are sketched in §5.2 and §5.3.

Acknowledgments. — The first author would like to thank Dominique Cerveau and Alcides Lins-Neto for pointing out and explaining [**10, 15**]. Both authors would like to thank Philippe Goutet for the pictures and for the nice presentation of the appendix.

2. Infinitesimal rigidity

2.1. Infinitesimal rigidity is a spectral property. —

Let X be any complex manifold. An *infinitesimal deformation* of X is an equivalence class of deformation at order one of a holomorphic cocycle defining X. These deformations are classified by the cohomology group $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ via the Kodaira-Spencer map.

Recall that in [**7**, §5] we have associated to any local holomorphic deformation $\mathfrak{X}$ a number of parameters $\mathfrak{m}(\mathfrak{X})$ that measures the variation of the complex structure of the generic fiber of $\mathfrak{X}$.

Definition 2.1. — An automorphism f of X is *rigid* if any deformation $\mathfrak{X}$ of X over a smooth base such that f lifts to an automorphism of $\mathfrak{X}$ satisfies $\mathfrak{m}(\mathfrak{X})=0$.

If X is a rational surface without holomorphic vector field, then $\mathfrak{X}$ satisfies $\mathfrak{m}(\mathfrak{X})=0$ if and only if $\mathfrak{X}$ is locally holomorphically trivial (because X admits a universal deformation over a smooth base).

Theorem 2.2. — *Let f be an automorphism of X and let f^* be the corresponding action of f on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$. If $\mathfrak{X}$ is a deformation of X such that f lifts to an automorphism of $\mathfrak{X}$, then*

$$\mathfrak{m}(\mathfrak{X})\leq \dim\ker\left(f^*-\mathrm{Id}\right).$$

In particular, if 1 *is not an eigenvalue of f^* then f is rigid.*

Proof. — Let b be a point in the base B of the deformation and let $\mathfrak{f}_b$ be the automorphism of $\mathfrak{X}$ lifting f. Since $\mathfrak{f}_b$ is a morphism of deformations, the Kodaira-Spencer map $\mathrm{KS}_b\colon \mathrm{T}_bB\longrightarrow \mathrm{H}^1(\mathfrak{X}_b,\mathrm{T}\mathfrak{X}_b)$ satisfies the identity $\mathfrak{f}_b^*\circ\mathrm{KS}_b=\mathrm{KS}_b$. Therefore the image of KS_b is contained in $\ker(\mathfrak{f}_b^*-\mathrm{Id})$. Let us recall that the rank of KS_b is equal to $\mathfrak{m}(\mathfrak{X})$ if b is generic. Thus, for generic b, we have the inequality $\mathfrak{m}(\mathfrak{X})\leq\dim\ker(\mathfrak{f}_b^*-\mathrm{Id})$. The function $b\mapsto\dim\ker(\mathfrak{f}_b^*-\mathrm{Id})$ being lower semi-continuous, we get that the inequality is valid for all b in B. □

We put the following definition:

Definition 2.3. — An automorphism f of X is *infinitesimally rigid* if f can not be lifted to an automorphism of a non-trivial infinitesimal deformation of X, or equivalently if 1 is not an eigenvalue of f^* acting on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$.

Note that an automorphism can be rigid without being infinitesimally rigid, but the converse holds: if f is infinitesimally rigid, then it is rigid.

2.2. Automorphisms of anticanonical basic rational surfaces. — *Anticanonical surfaces* are surfaces whose anticanonical class is effective. Let X be a basic rational anticanonical surface, and let D be an effective divisor representing the class $-\mathrm{K}_X$. Let us write X as a blowup

$$X=X_N\longrightarrow X_{N-1}\longrightarrow\ldots\longrightarrow X_1\longrightarrow X_0=\mathbb{P}^2,$$

and let $\pi_1,\ldots,\pi_N$ be the blowup maps and $E_1,\ldots,E_N$ be the exceptional divisors. Then there exists a sequence $(D_i)_{0\leq i\leq N}$ of effective divisors such that each D_i is an anticanonical divisor on X_i, $D_N=D$, and for any integer i between 1 and N, $D_i=\pi_i^*D_{i-1}-E_i$.

An anticanonical curve on $\mathbb{P}^2$ is a cubic. Then it is easy to prove that if D is reduced, then D is either an irreducible reduced curve of arithmetic genus 1, or a cycle of smooth rational curves, or one of the three exceptional configurations shown in the picture below:







If D is reducible but non reduced, the situation gets more complicated, even if we know that the irreducible components are still smooth rational curves. We start with the case of irreducible divisors.

Theorem 2.4. — *Let X be a basic rational surface with $\mathrm{K}_X^2 < 0$ endowed with an automorphism f, and assume that there exists an irreducible curve C on X which is linearly equivalent to $-\mathrm{K}_X$. Let P_f, Q_f and θ_C denote the characteristic polynomials of f^* acting on* $\mathrm{Pic}(X)$, $\mathrm{H}^1\big(X, \mathrm{T}X\big)$ *and* $\mathrm{H}^0\big(C, \mathrm{N}^*_{C/X}\big)$ *respectively.*

(a) If C is singular, $Q_f = P_f \times \theta_C$ and f is not infinitesimally rigid.

(b) If C is smooth, let us denote by ζ_C the action of f^* on the complex line $\mathrm{H}^0\big(C, \Omega^1_C\big)$.

Then $Q_f = P_f \times \theta_C \times \dfrac{x-\zeta_C}{x-1}$ and f^{12} is not infinitesimally rigid.

In both cases, X has no nonzero holomorphic vector field.

Proof. — Let us consider the long cohomology exact sequence associated with the short exact sequence

$$0 \longrightarrow \Omega^1_X(-C) \longrightarrow \Omega^1_X \longrightarrow \Omega^1_{X|C} \longrightarrow 0$$

of sheaves on X. Since $\mathrm{K}_X^2 < 0$ and since C is irreducible, $|-\mathrm{K}_X| = \{C\}$ so that f fixes globally the curve C. According to Serre duality for $0 \leq j \leq 2$,

$$\mathrm{H}^j\big(X, \mathrm{T}X\big)^* \simeq \mathrm{H}^{2-j}\big(X, \Omega^1_X \otimes \mathrm{K}_X\big) \simeq \mathrm{H}^{2-j}\big(X, \Omega^1_X(-C)\big).$$

On the other hand, by Hodge theory, $\mathrm{Pic}\,(X) \otimes_{\mathbb{Z}} \mathbb{C} \simeq \mathrm{H}^1\big(X, \Omega^1_X\big)$ and $\mathrm{H}^{2,0}(X) = \mathrm{H}^{0,2}(X) = 0$. Besides, $\mathrm{h}^2\big(X, \mathrm{T}X\big) = 0$ and $\mathrm{h}^2(X, \Omega^1_X) \simeq \mathrm{h}^0\big(X, \mathrm{T}X \otimes \mathrm{K}_X\big) = \mathrm{h}^0\big(X, \mathrm{T}X(-3)\big) = 0$. Hence we have an exact sequence of f-equivariant complex vector spaces

$$0 \longrightarrow \mathrm{H}^0\big(C, \Omega^1_{X|C}\big) \longrightarrow \mathrm{H}^1\big(X, \mathrm{T}X\big)^* \longrightarrow \mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C} \xrightarrow{\pi} \mathrm{H}^1\big(C, \Omega^1_{X|C}\big) \longrightarrow \mathrm{H}^0\big(X, \mathrm{T}X\big)^* \longrightarrow 0. \tag{2.1}$$

Even if C is singular, we have a conormal exact sequence

$$0 \longrightarrow \mathrm{N}^*_{C/X} \longrightarrow \Omega^1_{X|C} \longrightarrow \Omega^1_C \longrightarrow 0$$

where the injectivity of the first arrow holds because C is a local complete intersection in X. Since the arithmetic genus of C is 1, the dualizing sheaf ω_C is trivial. Besides, the conormal bundle $\mathrm{N}^*_{C/X}$ has degree $-\mathrm{K}_X^2$ which is positive, so that by Riemann-Roch and Serre duality, $\mathrm{h}^0\big(C, \mathrm{N}^*_{C/X}\big) = -\mathrm{K}_X^2$ and $\mathrm{h}^1\big(C, \mathrm{N}^*_{C/X}\big) = 0$. Therefore $\mathrm{H}^1\big(C, \Omega^1_{X|C}\big) \simeq \mathrm{H}^1\big(C, \Omega^1_C\big) \simeq \mathbb{C}$. It follows that π is the restriction morphism from $\mathrm{H}^1\big(X, \Omega^1_X\big)$ to $\mathrm{H}^1\big(C, \Omega^1_C\big)$.

Assume that C is singular. According to the adjunction formula,

$$\deg\, \Omega^1_C = \deg\, \Omega^1_{X|C} - \deg\, \mathrm{N}^*_{C/X} = \deg\, \Omega^2_{X|C} - \deg\, \mathrm{N}^*_{C/X} = \deg\, \omega_C = 0.$$

Besides, Ω^1_C is non-trivial since C is singular. Hence we obtain that $\mathrm{h}^0\big(C, \Omega^1_C\big)$ vanishes (otherwise we would have an exact sequence $0 \longrightarrow \mathcal{O}_C \longrightarrow \Omega^1_C \longrightarrow \mathcal{T} \longrightarrow 0$ where $\mathcal{T}$ is a torsion sheaf so that $\deg\, \Omega^1_C = \deg\, \mathcal{T}$ would be positive) so that by Riemann-Roch $\mathrm{h}^1\big(C, \Omega^1_C\big)$ vanishes as well. Therefore $\mathrm{h}^1\big(C, \Omega^1_{X|C}\big)$ vanishes, so that $\mathrm{h}^0\big(X, \mathrm{T}X\big) = 0$, and we obtain the f-equivariant exact sequence

$$0 \longrightarrow \mathrm{H}^0\big(C, \mathrm{N}^*_{C/X}\big) \longrightarrow \mathrm{H}^1\big(X, \mathrm{T}X\big)^* \longrightarrow \mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C} \longrightarrow 0$$

which gives $Q_f = P_f \times \theta_C$. The canonical class in $\mathrm{H}^1\big(X, \Omega^1_X\big)$ is invariant, 1 is thus always an eigenvalue of P_f, and an eigenvalue of Q_f as well.

Assume now that the curve C is smooth. Then $\mathrm{H}^1\big(C, \Omega^1_C\big)$ is canonically isomorphic to $\mathbb{C}$. Besides, the restriction morphism $\pi\colon \mathrm{H}^1\big(X, \Omega^1_X\big) \longrightarrow \mathrm{H}^1\big(C, \Omega^1_C\big) \simeq \mathbb{C}$ is surjective since $\pi([\mathrm{K}_X]) = \mathrm{K}_X.C = -C^2 > 0$. Therefore $\mathrm{h}^0\big(X, \mathrm{T}X\big) = 0$ and we have two f-equivariant exact sequences

$$0 \longrightarrow \mathrm{H}^0\big(C, \Omega^1_{X|C}\big) \longrightarrow \mathrm{H}^1\big(X, \mathrm{T}X\big)^* \longrightarrow \mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C} \longrightarrow \mathbb{C} \longrightarrow 0$$

$$0 \longrightarrow \mathrm{H}^0\big(C, \mathrm{N}^*_{C/X}\big) \longrightarrow \mathrm{H}^0\big(C, \Omega^1_{X|C}\big) \longrightarrow \mathrm{H}^0\big(C, \Omega^1_C\big) \longrightarrow 0.$$

This yields

$$Q_f \times (x-1) = P_f \times \theta_C \times (x-\zeta_C).$$

Therefore, ζ_C is an eigenvalue of Q_f. Since ζ_C belongs to the finite set $\{1, -1, \mathbf{i}, -\mathbf{i}, \mathbf{j}, \mathbf{j}^2\}$, we have $\zeta_C^{12} = 1$. Hence 1 is an eigenvalue of $Q_{f^{12}}$. □

We now investigate the case of anticanonical divisors given by a cycle of smooth rational curves and leave the three other exceptional configurations to the reader.

Remark that if f is an automorphism of a rational surface with positive entropy, then $|-\mathrm{K}_X|$ is either empty or consists of a single divisor (otherwise f would preserve a rational fibration). In this last case, by replacing f by an iterate, we can assume that all irreducible components of the divisor are globally invariant by f.

Theorem 2.5. — *Let X be a basic rational surface without nonzero holomorphic vector field and assume that there exists a reducible and reduced effective anticanonical cycle $D = \sum_{i=1}^r D_i$ on X such that $D_i^2 < 0$ for all i. For any automorphism f of X leaving each D_i globally invariant, let P_f, Q_f and θ_i denote the characteristic polynomials of f^* acting on* $\mathrm{Pic}(X)$, $\mathrm{H}^1\big(X, \mathrm{T}X\big)$ *and* $\mathrm{H}^0\big(D_i, \mathrm{N}^*_{D_i/X}(-S_i)\big)$ *respectively, where $S_i = \mathrm{sing}(D) \cap D_i$. Then*

$$Q_f = P_f \times \frac{\prod_{i=1}^r \theta_i}{(x-1)^r}.$$

Proof. — Let S be the singular locus of D. Then we have an exact sequence

$$0 \longrightarrow \Omega^1_{X|D} \longrightarrow \oplus_{i=1}^r \Omega^1_{X|D_i} \longrightarrow \Omega^1_{X|S} \longrightarrow 0.$$

Since $N^*_{D_i/X}$ has positive degree, $\mathrm{H}^1\big(D_i, \Omega^1_{X|D_i}\big) \simeq \mathrm{H}^1\big(D_i, \Omega^1_{D_i}\big) \simeq \mathbb{C}$. Besides, since $\mathrm{H}^0\big(D_i, \Omega^1_{D_i}\big) = 0$, we obtain the isomorphism $\mathrm{H}^0\big(D_i, \Omega^1_{X|D_i}\big) \simeq \mathrm{H}^0\big(D_i, N^*_{D_i/X}\big)$. It follows that

$$\mathrm{H}^0\big(D, \Omega^1_{X|D}\big) \simeq \bigoplus_{i=1}^r \mathrm{H}^0\big(D, N^*_{D_i/X}(-S_i)\big).$$

and that the map from $\bigoplus_{i=1}^r \mathrm{H}^0\big(D_i, \Omega^1_{X|D_i}\big)$ to $\mathrm{H}^0\big(S, \Omega^1_{X|S}\big)$ is onto. Therefore

$$\mathrm{H}^1\big(D, \Omega^1_{X|D}\big) \simeq \bigoplus_{i=1}^r \mathrm{H}^1\big(D_i, \Omega^1_{X|D_i}\big) \simeq \bigoplus_{i=1}^r \mathrm{H}^1\big(D_i, \Omega^1_{D_i}\big) \simeq \mathbb{C}^r.$$

We now write down the f-equivariant exact sequence (2.1) used in the proof of Theorem 2.4:

$$0 \longrightarrow \bigoplus_{i=1}^r \mathrm{H}^0\big(D, N^*_{D_i/X}(-S_i)\big) \longrightarrow \mathrm{H}^1\big(X, \mathrm{T}X\big)^* \longrightarrow \mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C} \longrightarrow \mathbb{C}^r \longrightarrow \mathrm{H}^0\big(X, \mathrm{T}X\big)^* \longrightarrow 0.$$

Since we assumed that there were no nonzero holomorphic vector fields on X, we get $Q_f \times (x-1)^r = P_f \times \prod_{i=1}^r \theta_i$. □

3. A geometric example: Kummer surfaces

We start by fixing some conventions and notations. For any group G and any vector space W, a G-module means a *contravariant* representation of G in W. For any element g in a group G, we write $\langle g \rangle$ for the cyclic group in G generated by the element g. We denote the vector space $\mathbb{C}^2$ by V. If nothing else is specified, we consider V as a natural $\mathrm{GL}(2;\Lambda)$-module, where a matrix M acts by M^{-1}. Except explicit mention of the contrary, the letter Λ always denotes the hexagonal lattice $\mathbb{Z}[\mathbf{j}]$ in $\mathbb{C}^2$, where $\mathbf{j} = e^{2\mathbf{i}\pi/3}$.

3.1. Construction, Picard group and fixed points. — Let $\mathcal{E}$ be the elliptic curve obtained by taking the quotient of the complex line $\mathbb{C}$ by the hexagonal lattice Λ and let A be the abelian surface $\mathcal{E} \times \mathcal{E}$. The group $\mathrm{GL}(2;\Lambda)$ acts linearly on $\mathbb{C}^2$ and preserves the lattice $\Lambda \times \Lambda$. Therefore any element M of $\mathrm{GL}(2;\Lambda)$ induces an automorphism f_M on A that commutes with the automorphism ϕ of A of order three defined by $\phi(x,y) = (\mathbf{j}x, \mathbf{j}y)$. It follows that f_M induces an automorphism of the singular surface $A/\langle\phi\rangle$. After blowing up the singular points, we obtain an automorphism $\widetilde{f_M}$ on a smooth model X, which is a rational Kummer surface.

Let us remark that ϕ and ϕ^2 have the same 9 fixed points (we will see that these points also appear in §3.2):

$$p_1=(0,0),\, p_2=\left(0,\frac{2}{3}+\frac{\mathbf{j}}{3}\right),\, p_3=\left(0,\frac{1}{3}+\frac{2\mathbf{j}}{3}\right),\, p_4=\left(\frac{2}{3}+\frac{\mathbf{j}}{3},0\right),\, p_5=\left(\frac{2}{3}+\frac{\mathbf{j}}{3},\frac{2}{3}+\frac{\mathbf{j}}{3}\right),$$

$$p_6=\left(\frac{2}{3}+\frac{\mathbf{j}}{3},\frac{1}{3}+\frac{2\mathbf{j}}{3}\right),\, p_7=\left(\frac{1}{3}+\frac{2\mathbf{j}}{3},0\right),\, p_8=\left(\frac{1}{3}+\frac{2\mathbf{j}}{3},\frac{2}{3}+\frac{\mathbf{j}}{3}\right),\, p_9=\left(\frac{1}{3}+\frac{2\mathbf{j}}{3},\frac{1}{3}+\frac{2\mathbf{j}}{3}\right).$$

In particular the group $G=\langle\phi\rangle\simeq\mathbb{Z}_3$ has 9 fixed points on A. We denote this finite set by S, it is a subgroup of the torsion points of A of order 3. Note that the group $\mathrm{GL}(2;\Lambda)$ leaves the set S globally invariant. We denote by H the kernel of the natural group morphism from $\mathrm{SL}(2;\Lambda)$ to $\mathrm{SL}(2;\mathbb{F}_9)$ where $\mathbb{F}_9=\mathbb{Z}_3[\mathbf{j}]$. This is a subgroup of finite index in $\mathrm{GL}(2;\Lambda)$ which fixes every point in S.

There is a slightly different way to construct X: first we blow up the set S in $A=\mathcal{E}\times\mathcal{E}$ and denote by $\widetilde{A}$ the resulting surface and by $\widetilde{E}_i$ the exceptional divisors corresponding to the points p_i. The group G acts on $\widetilde{A}$, and the quotient $\widetilde{A}/G$ is exactly our Kummer surface X. Then we have the following diagram, where δ is the blowup map and π denotes the projection map from $\widetilde{A}$ to X:

$$\begin{tikzcd} & \widetilde{A} \arrow[dl, "\delta"'] \arrow[dr, "\pi"] & \\ A & & X \end{tikzcd}$$

This diagram is equivariant for both actions of G and $\mathrm{GL}(2;\Lambda)$. We set $E_i=\pi(\widetilde{E}_i)$. Then a straightforward computation shows that π is the projection of the cyclic cover of X of order 3 along each rational curve E_i.

Let us recall that X is obtained by blowing up the projective plane $\mathbb{P}^2$ in 12 distinct points. The number of blown up points can be determined easily by computing the Euler characteristic of X: indeed, $\chi(\widetilde{A})=\chi(\mathrm{Bl}_S A)=\chi(A)+9=9$. As $\#G=3$, we get $\chi\big(\widetilde{A}\smallsetminus\{\widetilde{E}_i\}_i\big)=3\chi\big(X\smallsetminus\{E_i\}_i\big)$. But $\chi\big(\widetilde{A}\smallsetminus\{\widetilde{E}_i\}_i\big)=-9$ so $\chi\big(X\smallsetminus\{E_i\}_i\big)=-3$ and

$$\chi(X)=-3+18=15=\chi\big(\mathbb{P}^2\big)+12.$$

Another way to see it is given in §3.2.

The Picard group of X can be described in the following way: since X is rational, $\mathrm{Pic}(X)\simeq\mathrm{H}^2(X,\mathbb{Z}_X)$. First we compute:

$$\mathrm{H}^2\big(\widetilde{A},\mathbb{Z}_{\widetilde{A}}\big)\simeq\mathrm{H}^2(A,\mathbb{Z}_A)\oplus\left(\bigoplus_{i=1}^{9}\mathbb{Z}[\widetilde{E}_i]\right)\simeq\wedge^2_{\mathbb{Z}}(\Lambda^*\times\Lambda^*)\oplus\left(\bigoplus_{i=1}^{9}\mathbb{Z}[\widetilde{E}_i]\right),$$

where Λ^* denotes the dual lattice of Λ. Then we get an isomorphism of $\mathrm{GL}(2;\Lambda)$-modules:

$$\mathrm{Pic}(X)\simeq\mathrm{H}^2\big(\widetilde{A},\mathbb{Z}_{\widetilde{A}}\big)^G\simeq\big(\wedge^2_{\mathbb{Z}}(\Lambda^*\times\Lambda^*)\big)^G\oplus\bigoplus_{i=1}^{9}\mathbb{Z}[\widetilde{E}_i]. \tag{3.1}$$

It is easy to see that $\big(\wedge^2_{\mathbb{Z}}(\Lambda^*\times\Lambda^*)\big)^G$ is a free $\mathbb{Z}$-module of rank 4, so that $\mathrm{Pic}(X)$ is a free $\mathbb{Z}$-module of rank 13 (which is the number of blows up plus 1, as expected).

Let M be $\mathrm{GL}(2;\Lambda)$ with eigenvalues α and β. Then the complex eigenvalues of the endomorphism $\wedge^2M$ of the real vector space $\wedge^2_{\mathbb{R}}V$ are

$$|\alpha|^2,\qquad|\beta|^2,\qquad\alpha\overline{\beta},\qquad\overline{\alpha}\beta,\qquad\alpha\beta\qquad\text{and}\qquad\overline{\alpha}\overline{\beta}.$$

A straightforward but lengthy computation shows that the eigenvalues of $\wedge^2M$ on the subspace $\big(\wedge^2_{\mathbb{R}}V\big)^G$ are exactly $|\alpha|^2$, $|\beta|^2$, $\alpha\overline{\beta}$ and $\overline{\alpha}\beta$. Thus, for any element M in $\mathrm{GL}(2;\Lambda)$ with spectral radius r_M, the spectral radius of M acting on $\big(\wedge^2_{\mathbb{Z}}(\Lambda^* \times \Lambda^*)\big)^G$ is r_M^2. This means that the first dynamical degree (1) of $\widetilde{f}_M$ is given by the formula $\lambda_1(\widetilde{f}_M) = r_M^2$. More precisely, the characteristic polynomial of $\widetilde{f}_M^*$ acting on $\mathrm{Pic}(X)$ is

$$(x-1)^9\big(x-|\alpha|^2\big)\big(x-|\beta|^2\big)\big(x-\overline{\alpha}\beta\big)\big(x-\alpha\overline{\beta}\big).$$

In the remaining part of this section, we compute the number of fixed points of $\widetilde{f}_M$ for any M in H. As a consequence, we compute the trace of the endomorphism $\widetilde{f}_M^*$ with the help of Atiyah-Bott fixed point formula. We will obtain a much finer result in the next section (Theorem 3.7).

In the sequel, we divide the set of fixed points of $\widetilde{f}_M$ into two parts: the first part Θ_1 consists of fixed points outside the exceptional divisors E_i; and the second part Θ_2 consists of the remaining fixed points.

Proposition 3.1. — *Let M be a matrix in H such that $M^3 \neq \mathrm{Id}$. Then the automorphism $\widetilde{f}_M$ has $|\mathrm{Tr}(M)|^2 + 11$ fixed points on the Kummer surface X.*

Proof. — The first step consists in proving that the fixed points of $\widetilde{f}_M$ are non-degenerate, *i.e.* that 1 is never an eigenvalue of the differential of $\widetilde{f}_M$ at a fixed point. We deal with fixed points in Θ_1 and Θ_2 separately.

The points of Θ_1 correspond to points p in $A \setminus S$ such that $f_M(p)$ lies in the orbit $G \cdot p$, modulo the action of G. For any such point, the differential of $\widetilde{f}_M$ identifies with M, $\mathbf{j}M$ or $\mathbf{j}^2M$, so that 1 is never an eigenvalue.

The set Θ_2 can be described as follows: the automorphism $\widetilde{f}_M$ acts by the projective transformation $\mathbb{P}(M)$ on the curve E_i via the identification

$$E_i \simeq \widetilde{E}_i \simeq \mathbb{P}(\mathrm{T}_{p_i}V) \simeq \mathbb{P}(V).$$

Therefore we get two fixed points q_α and q_β on each E_i corresponding to the eigenspaces of M. This gives a concrete description of Θ_2, which consists of 18 points (two distinct points on each E_i). To compute the differential of $\widetilde{f}_M$ at such a fixed point, we can assume without loss of generality that this point lies in E_1. Let (e,f) be an eigenvector of M for the eigenvalue α and assume for simplicity that $e \neq 0$. If (x,y) are the standard coordinates on V, we define coordinates (u,v) in $\widetilde{A}$ near $\widetilde{E}_1$ by putting $x = u$ and $y = uv$. Then if we set $Z = u^3$ and $T = v$, (Z,T) are holomorphic coordinates on X near q_α. In these coordinates, $q_\alpha = \left(0, \dfrac{f}{e}\right)$ and $\widetilde{f}_M$ is given by

$$(Z,T) \mapsto \left(Z(a+bT)^3, \frac{c+dT}{a+bT}\right).$$

so that the eigenvalues of $\mathrm{d}\widetilde{f}_M$ at q_α are α^3 and α^{-2}, which are different from 1.

We can now apply the Lefschetz fixed point formula. As $\mathrm{H}^1\big(X,\mathbb{R}\big) = \mathrm{H}^3\big(X,\mathbb{R}\big) = 0$, we get that the number of fixed points of $\widetilde{f}_M$ is equal to $\mathrm{Tr}\, \widetilde{f}^{\,*}_{M\,|\mathrm{H}^2(X,\mathbb{R})} + 2$. Then we can use formula (3.1): the trace on the part $\big(\wedge^2_{\mathbb{Z}}(\Lambda^* \times \Lambda^*)\big)^G$ is the sum

$$|\alpha|^2 + |\beta|^2 + \alpha\overline{\beta} + \overline{\alpha}\beta = |\alpha+\beta|^2;$$

and since M belongs to H, $\widetilde{f}_M$ acts trivially on the factor $\bigoplus_{i=1}^9 \mathbb{Z}[\widetilde{E}_i]$ so that the trace on this factor is 9. This gives the required result. □

We will now study in greater details the set Θ_1. Let us define three subsets S_1, S_2, S_3 of A as follows:

$$S_1 = \{p \in A \setminus S \mid f_M(p) = p\}, \qquad S_2 = \{p \in A \setminus S \mid f_M(p) = \phi(p)\}, \qquad S_3 = \{p \in A \setminus S \mid f_M(p) = \phi^2(p)\}.$$

To compute the cardinality of the S_i's, we use the following statement:

Lemma 3.2. — *For any matrix P in $\mathrm{GL}(2;\Lambda)$ such that 1 is not an eigenvalue of P, the automorphism f_P of the complex torus A has $\big|1 - \mathrm{Tr}(P) + \det P\big|^2$ fixed points.*

---

1. Let f be a bimeromorphic map on a Kähler surface X. The *first dynamical degree* of f is given by $\lambda_1(f) = \lim_{n\to+\infty}(|(f^n)^*|)^{1/n}$ where f^* is the induced map from $\mathrm{H}^{1,1}\big(X,\mathbb{R}\big)$ into itself.

Proof. — Since the fixed points of f_P are non-degenerate, we can apply the holomorphic Lefschetz fixed point formula [**9**, p. 426]; this gives

$$1-\overline{\mathrm{Tr}\,(P)}+\overline{\det P}=\#\mathrm{Fix}\,(f_P)\times\frac{1}{1-\mathrm{Tr}\,(P)+\det P}.$$

□

Corollary 3.3. — *Let M be a matrix in H such that $M^3\neq\mathrm{Id}$. One has the following equalities:*

$$\#S_1=\left|\mathrm{Tr}\,(M)-2\right|^2-9\qquad\textit{and}\qquad\#S_2=\#S_3=\left|1+\mathrm{Tr}\,(M)\right|^2-9.$$

Proof. — This follows directly from Lemma 3.2, since

$$S_1=\mathrm{Fix}\,(f_M)\smallsetminus S,\qquad S_2=\mathrm{Fix}\,(f_{\mathbf{j}^2M})\smallsetminus S\qquad\text{and}\qquad S_3=\mathrm{Fix}\,(f_{\mathbf{j}M})\smallsetminus S.$$

□

The group G acts transitively on each set S_i. Then Θ_1 can be written as the disjoint union of the quotients S_i/G. Remark that we can obtain in this way the result of Proposition 3.1: indeed, if $t=\mathrm{Tr}\,(M)$,

$$\#\Theta_1=\frac{|t-2|^2}{3}-3+2\left(\frac{|1+t|^2}{3}-3\right)=|t|^2-7$$

Since $\#\Theta_2=18$, this gives $\#\mathrm{Fix}\,(\widetilde{f}_M)=|t|^2+11$.

As $\mathrm{H}^0\big(X,\mathrm{T}X\big)=\mathrm{H}^2\big(X,\mathrm{T}X\big)=0$, the holomorphic Atiyah-Bott formula [**1**, Thm. 4.12] yields the following result:

Proposition 3.4. — *The trace of $\widetilde{f}_M^{\,*}$ acting on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ is equal to the sum $-\sum_x\dfrac{\mathrm{Tr}\ \mathrm{d}f_x^{-1}}{\det(\mathrm{Id}-\mathrm{d}f_x)}$ where x runs through the fixed points of $\widetilde{f}_M$.*

Corollary 3.5. — *Let M be a matrix in H such that $M^3\neq\mathrm{Id}$. Then the trace of $\widetilde{f}_M^{\,*}$ acting on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ is $8\,\mathrm{Tr}\,(M)$.*

Proof. — We divide the fixed point set of $\widetilde{f}_M$ in four parts: S_1/G, S_2/G, S_3/G that form Θ_1 and Θ_2. We start by computing the contribution of Θ_2 in the sum of Proposition 3.4. Let us put $t=\mathrm{Tr}\,(M)$. Any pair of fixed points in a divisor E_i yields the term

$$\frac{\alpha^{-3}+\alpha^2}{(1-\alpha^3)(1-\alpha^{-2})}+\frac{\alpha^3+\alpha^{-2}}{(1-\alpha^{-3})(1-\alpha^2)}$$

so that the contribution of Θ_2 is $-9t-\dfrac{6}{t-2}-\dfrac{3}{1+t}$. Now the contributions of S_1/G, S_2/G and S_3/G are respectively

$$\left(\frac{|t-2|^2}{3}-3\right)\frac{t}{2-t},\qquad\left(\frac{|1+t|^2}{3}-3\right)\frac{-\mathbf{j}^2t}{1+t}\qquad\text{and}\qquad\left(\frac{|1+t|^2}{3}-3\right)\frac{-\mathbf{j}t}{1+t}$$

so that the contribution of Θ_1 is $t+\dfrac{3t}{t-2}-\dfrac{3t}{1+t}$. Adding the contributions of Θ_1 and Θ_2 we get $-8t$. □

3.2. An alternative construction via foliations. — In [**10, 15**] the authors consider the pencil $\mathcal{P}_4 = \{\mathcal{F}_\alpha\}_{\alpha\in\mathbb{P}^1}$ in $\mathbb{P}^2$ where $\mathcal{F}_\alpha$ is given by the 1-form $\omega + \alpha\eta$ with

$$\omega = \big(x^3-1\big)x\,dy - \big(y^3-1\big)y\,dx, \qquad \eta = \big(x^3-1\big)x\,dy - \big(y^3-1\big)x^2\,dx.$$

The tangency set of $\mathcal{P}_4$, given by $\omega\wedge\eta = 0$, is an algebraic curve; more precisely it is the union of 9 invariant lines that intersect at 12 points. Let $\mathcal{C}$ be the union of these points and lines. Let (x,y) be a system of coordinates of $\mathbb{C}^2$ and let $\{F_\alpha\}_\alpha$ be the pencil of linear foliations on $\mathbb{C}^2$ where F_α is described by $\Omega_\alpha = dy - \alpha dx$. Let us denote by $p\colon\, \mathbb{C}^2 \longrightarrow A$ the canonical projection and by $\mathcal{Q}_4$ the pencil of foliations $\{\mathcal{G}_\alpha\}_\alpha = \{p_*F_\alpha\}_\alpha$. Let us consider the following elliptic curves on A:

$$\mathcal{E}_{0,1} = \{0\}\times\mathcal{E}, \qquad \mathcal{E}_{1,0} = \mathcal{E}\times\{0\}, \qquad \mathcal{E}_{1,1} = \big\{(x,x)\,|\, x\in\mathcal{E}\big\}, \qquad \mathcal{E}_{1,\mathbf{j}} = \big\{(x,-\mathbf{j}x)\,|\,x\in\mathcal{E})\big\}.$$

Set $\mathscr{C} = \big\{\mathcal{E}_{0,1},\, \mathcal{E}_{1,0},\, \mathcal{E}_{1,1},\, \mathcal{E}_{1,\mathbf{j}}\big\}$. Given F in $\mathscr{C}$ and m in $\mathrm{Fix}(\phi)$ denote $F_m = F + m$. The set

$$\mathscr{E} = \big\{F_m\,|\,m\in\mathrm{Fix}(\phi),\, F\in\mathscr{C}\big\}$$

consists in 12 elliptic curves $\mathcal{E}_1$, $\ldots$, $\mathcal{E}_{12}$; two such different elliptic curves intersect only at 3 fixed points of $\mathrm{Fix}(\phi)$. Let $\mathcal{C}^*$ be the union of $\mathrm{Fix}(\phi)$ and these 12 elliptic curves.

According to McQuillan [**5**, Chapter 8, Theorem 1] there exists a rational map $\psi\colon A \dashrightarrow \mathbb{P}^2$ such that $\psi^*(\mathcal{P}_4) = \mathcal{Q}_4$. Let us explain the construction of ψ. Since the restriction of the morphism π of degree 3 to $\widetilde{E}_i$ is a biholomorphism, π maps any $\widetilde{E}_i$ to a rational curve of self-intersection -3. Besides π maps each elliptic curve $\pi^*\mathcal{E}_i$, $\mathcal{E}_i\in\mathscr{E}$, to a rational curve $\widetilde{\mathcal{E}}_i$ with self-intersection -1. Denote by $\pi_1 : X \to X_0$ the blowing-down map of the curves $\widetilde{\mathcal{E}}_1$, $\ldots$, $\widetilde{\mathcal{E}}_{12}$; one can see that $X_0 = \mathbb{P}^2$ (*see* [**15**, Lemma 3]). The map ψ is given by

$$\psi = \pi_1\pi\delta^{-1} : A \dashrightarrow \mathbb{P}^2.$$

Set $\mathscr{E}_* = \psi(\mathscr{E})$, $\mathrm{Fix}(\phi)_* = \psi(\mathrm{Fix}(\phi))$ and let $\mathcal{C}_*$ be the union of $\mathscr{E}_*$ and $\mathrm{Fix}(\phi)_*$. In [**10**, Proposition 1] the author proves that $\mathcal{C} = \mathcal{C}_*$ up to an automorphism of $\mathbb{P}^2$. In particular X is obtained by blowing up $\mathbb{P}^2$ in 12 points.

The family $\{\mathcal{F}_\alpha\}_\alpha$ plays a very important role in the study of foliations; it allows to give a negative answer to the two famous following problems [**10**]:

- Poincaré Problem: "is it possible to decide if an algebraic differential equation in two variables is algebraically integrable (in the sense that it has a rational first integral) ?" [**12, 14, 13**]
- "Is it possible to recognize the genus of the general solution of an algebraic differential equation in two variables which has a rational first integral ?"

Let M be an element of $\mathrm{GL}(2;\Lambda)$ with two eigenvalues μ_1, μ_2 such that $|\mu_1| < 1 < |\mu_2|$. The linear transformation induced by M preserves the pencil $\{F_\alpha\}_\alpha$ and fixes F_{β_1/γ_1}, F_{β_2/γ_2} where (β_1,γ_1) and (β_2,γ_2) are the two eigenvectors of ${}^tM$; as a consequence $\widetilde{f}_M$ preserves $\pi^*\delta^*p_*F_{\beta_1/\gamma_1}$ and $\pi^*\delta^*p_*F_{\beta_2/\gamma_2}$.

3.3. Action of the automorphism group on infinitesimal deformations. — In this section we study the $\mathrm{GL}(2;\Lambda)$-module $\mathrm{H}^1(X,\mathrm{T}X)$ where X is the Kummer surface associated to an hexagonal lattice (we also give the results for cubical lattices at the end of the section). For this we will use two fundamental exact sequences:

First exact sequence relating $\mathrm{T}A$ *and* $\mathrm{T}\widetilde{A}$ *via the blowup map* δ:

$$0 \longrightarrow \mathrm{T}\widetilde{A} \longrightarrow \delta^*\mathrm{T}A \longrightarrow \bigoplus_{i=1}^{9}\iota_{E_i*}\mathrm{T}_{\widetilde{E}_i}(-1) \longrightarrow 0 \tag{3.2}$$

This sequence is $\mathrm{GL}(2;\Lambda)$-equivariant and G-equivariant. Remark that $\mathrm{T}_{\widetilde{E}_i}(-1)$ is isomorphic to $\mathcal{O}_{E_i}(1)$, but this isomorphism is not canonical and cannot be made compatible in any way with the action of $\mathrm{GL}(2;\Lambda)$.

Second exact sequence relating $\mathrm{T}\widetilde{A}$ *and* $\mathrm{T}X$ *via the cyclic cover* π:

$$0 \longrightarrow \big(\pi_*\mathrm{T}\widetilde{A}\big)^G \longrightarrow \mathrm{T}X \longrightarrow \bigoplus_{i=1}^{9}\iota_{E_i*}\mathrm{N}_{E_i/X} \longrightarrow 0 \tag{3.3}$$

This sequence is $\mathrm{GL}(2;\Lambda)$-equivariant.

Let us denote by $\mathcal{F}$ the sheaf $\displaystyle\bigoplus_{i=1}^{9} \iota_{E_i *}\mathrm{T}_{\widetilde{E}_i}(-1)$. We state and prove all technical results we need.

Lemma 3.6. — *The following assertions hold :*

(a) For any integer i with $0 \leq i \leq 2$, the natural morphisms

$$\begin{cases} \mathrm{H}^i\big(A,\mathrm{T}A\big) \longrightarrow \mathrm{H}^i\big(\widetilde{A},\delta^*\mathrm{T}A\big) \\ \mathrm{H}^i\big(X,\pi_*\mathrm{T}\widetilde{A}\big) \longrightarrow \mathrm{H}^i\big(\widetilde{A},\pi^*\pi_*\mathrm{T}\widetilde{A}\big) \longrightarrow \mathrm{H}^i\big(\widetilde{A},\mathrm{T}\widetilde{A}\big) \end{cases}$$

are $\mathrm{GL}(2;\Lambda)$-equivariant isomorphisms.

(b) The cohomology groups $\mathrm{H}^0\big(\widetilde{A},\mathcal{F}\big)^G$ and $\mathrm{H}^1\big(\widetilde{A},\mathcal{F}\big)^G$ vanish.

(c) The cohomology group $\mathrm{H}^1\big(X,(\pi_*\mathrm{T}\widetilde{A})^G\big)$ vanishes. Besides, $\mathrm{H}^2\big(X,(\pi_*\mathrm{T}\widetilde{A})^G\big)$ is isomorphic to V as an H-module.

(d) For $1 \leq i \leq 9$, we have an isomorphism of H-modules between $\mathrm{H}^1\big(E_i,\mathrm{N}_{E_i/X}\big)$ and V.

Proof. — *(a)* We write the Leray spectral sequence for the pair $(\delta^*\mathrm{T}A, A)$: we have

$$\mathrm{E}_2^{p,q} = \mathrm{H}^p\big(A,\mathrm{R}^q\delta_*\mathcal{O}_{\widetilde{A}}\otimes\mathrm{T}A\big) \qquad \text{and} \qquad \mathrm{E}_\infty^{p,q} = \mathrm{Gr}^p\mathrm{H}^{p+q}\big(\widetilde{A},\delta^*\mathrm{T}A\big).$$

Since δ is the projection of a point blowup, $\mathrm{R}^q\delta_*\mathcal{O}_{\widetilde{A}} = 0$ for $q > 0$ so that the spectral sequence degenerates and we get $\mathrm{H}^p(A,\mathrm{T}A) = \mathrm{E}_2^{p,0} \simeq \mathrm{E}_\infty^{p,0} = \mathrm{H}^p\big(\widetilde{A},\delta^*\mathrm{T}A\big)$. The argument is the same for the second morphism: π being finite, $\mathrm{R}^q\pi_*\mathrm{T}\widetilde{A} = 0$ for $q > 0$.

(b) The vanishing of $\mathrm{H}^1\big(\widetilde{A},\mathcal{F}\big)^G$ is straightforward as $\mathrm{H}^1(\widetilde{A},\mathcal{F}) = 0$. We see that $\mathrm{H}^0(\widetilde{A},\mathcal{F})$ is isomorphic as a G-module to the direct sum $\bigoplus_{i=1}^{9}\mathrm{T}_{p_i}A$, where ϕ acts by the inverse of its differential at each point p_i. Therefore, $\mathrm{H}^0(\widetilde{A},\mathcal{F}) \simeq V^9$ and the result follows since $V^G = \{0\}$.

(c) According to *(a)* and since G is finite, we have isomorphisms

$$\mathrm{H}^i\big(X,(\pi_*\mathrm{T}\widetilde{A})^G\big) \simeq \mathrm{H}^i\big(X,(\pi_*\mathrm{T}\widetilde{A})\big)^G \simeq \mathrm{H}^i\big(\widetilde{A},\mathrm{T}\widetilde{A}\big)^G.$$

Using (3.2) combined with *(b)* and *(a)*, we obtain $\mathrm{GL}(2;\Lambda)$-equivariant isomorphisms $\mathrm{H}^i\big(\widetilde{A},\mathrm{T}\widetilde{A}\big)^G \simeq \mathrm{H}^i\big(A,\mathrm{T}A\big)^G$ for $i = 1,2$. Now $\mathrm{H}^i\big(A,\mathrm{T}A\big)^G$ is isomorphic to $V^*\otimes V$ (resp. $\wedge^2 V^*\otimes V$) for $i = 1$ (resp. $i = 2$), where a matrix M in $\mathrm{GL}(2;\Lambda)$ acts by ${}^t\overline{M}\otimes M^{-1}$ $\big($resp. $\wedge^2\,{}^t\overline{M}\otimes M^{-1}\big)$. This proves that $\mathrm{H}^1\big(A,\mathrm{T}A\big)^G$ vanishes and that $\mathrm{H}^2\big(A,\mathrm{T}A\big)^G = \mathrm{H}^2\big(A,\mathrm{T}A\big)$ since $\overline{\mathbf{j}}^2\,\mathbf{j}^{-1} = 1$. This yields the result.

(d) If $W = \mathrm{T}_{p_i}A$, we have the Euler exact sequence on E_i :

$$0 \longrightarrow \mathcal{O}_{E_i}(-1) \longrightarrow W\otimes\mathcal{O}_{E_i} \longrightarrow \mathrm{T}E_i(-1) \longrightarrow 0$$

which proves that the canonical bundle of E_i is canonically isomorphic to $\det W^*\otimes_{\mathbb{C}}\mathcal{O}_{E_i}(-2)$. Since $\mathrm{N}_{E_i/X}$ is canonically isomorphic to $\mathcal{O}_{E_i}(-3)$, we get by Serre duality a chain of H-equivariant isomorphisms

$$\mathrm{H}^1\big(E_i,\mathrm{N}_{E_i/X}\big) \simeq \mathrm{H}^0\big(E_i,\mathcal{O}_{E_i}(3)\otimes\mathrm{K}_{E_i}\big)^* \simeq \det W\otimes\mathrm{H}^0\big(E_i,\mathcal{O}_{E_i}(1)\big)^* \simeq \det W\otimes W \simeq \det V\otimes V.$$

This finishes the proof of the statement, since $\det V$ is trivial as an H-module. ☐

Theorem 3.7. — *For any element M in H, the characteristic polynomial of the endomorphism $\widetilde{f}_M^{\,*}$ of $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ is equal to $\big(x^2-\mathrm{tr}(\mathrm{M})\,x+1\big)^8$.*

Proof. — Using the exact sequence (3.3), Lemma 3.6 *(c)*, *(d)* and the fact that $\mathrm{H}^2\big(X,\mathrm{T}X\big) = 0$, we get an exact sequence of H-modules

$$0 \longrightarrow \mathrm{H}^1\big(X,\mathrm{T}X\big) \longrightarrow V^9 \longrightarrow V \longrightarrow 0.$$

The result follows since the characteristic polynomial of an endomorphism is multiplicative under equivariant short exact sequences. ☐

Corollary 3.8. — *For any matrix M of infinite order in* $\mathrm{GL}(2;\Lambda)$*, $\widetilde{f}_M$ is a rigid automorphism.*

Proof. — Let M be a matrix of infinite order in $\mathrm{GL}(2;\Lambda)$. Since H has finite index in $\mathrm{GL}(2;\Lambda)$, there exists an integer p such that M^p lies in H. By Theorem 3.7, 1 is not an eigenvalue of $\widetilde{f}^*_{M^p}$ $\big($and therefore of $\widetilde{f}^*_M\big)$. Thus $\widetilde{f}_M$ is infinitesimally rigid so it is rigid. □

For the sake of completeness, we also provide briefly the corresponding results for cubical lattices. The same strategy works but the results are completely different. In the cubical case, the group G is of order 4, the map ϕ (which is this case the multiplication by $\mathbf{i}$) has 4 fixed points and 12 new other points are fixed by ϕ^2. After blowing up these 16 points and taking the quotient by G, we get a rational Kummer surface X that can also be obtained by blowing up $\mathbb{P}^2$ in 13 points. Besides, $\widetilde{A}$ is the cyclic covering of X along 10 disjoint rational curves $E_1,\ldots,E_{10}$, where the multiplicity of the covering is 4 along E_1, E_2, E_3, E_4 and 2 along the other ten remaining E_i's. We can define the subgroup K of finite index in $\mathrm{SL}(2;\Lambda)$ consisting of the elements fixing the 16 points. In the cubic case $\mathrm{H}^0\big(X,\pi_*\mathrm{T}\widetilde{A}\big)^G=\mathrm{H}^1\big(X,\pi_*\mathrm{T}\widetilde{A}\big)^G=0$. Therefore we get an isomorphism of K-modules

$$\mathrm{H}^1\big(X,\mathrm{T}X\big)\simeq\bigoplus_{i=1}^{10}\mathrm{H}^1\big(E_i,\mathrm{N}_{E_i/X}\big).$$

The first four terms correspond to the K-module $\mathrm{Sym}^2\,\mathrm{T}_{p_i}A$ (which is of rank 3) and the six last ones to the trivial K-module $\mathbb{C}$. Therefore we get

Theorem 3.9. — *For a cubic Kummer surface X the K-module* $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ *of infinitesimal deformations is isomorphic to* $\big(\mathrm{Sym}^2 V\big)^4\oplus\mathbb{C}^6$*. In particular, for any element M in K, the characteristic polynomial of the endomorphism $\widetilde{f}^*_M$ of* $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ *is equal to* $\Big(\chi_{\mathrm{Sym}^2 M^{-1}}(x)\Big)^4(x-1)^6$.

As a corollary, we see that elements in K are not infinitesimally rigid. We can prove a slightly stronger result: since $(0,0)$ is always fixed by any matrix M, $\mathrm{H}^1\big(E_1,\mathrm{N}_{E_1/X}\big)$ is a $\mathrm{GL}(2;\Lambda)$-submodule of $\mathrm{H}^1\big(X,\mathrm{T}X\big)$, it is isomorphic to $\det V\otimes\mathrm{Sym}^2 V$. Therefore, $\widetilde{f}_M$ is not infinitesimally rigid as soon as $(\det M)^2=1$. Nevertheless it is possible to construct infinitesimally rigid elements in $\mathrm{GL}(2;\Lambda)$, but they are of lower interest than in the hexagonal case because they do not remain infinitesimally rigid after iteration. For instance, we have the following result:

Theorem 3.10. — *Let M be a matrix in* $\mathrm{GL}(2;\Lambda)$ *such that* $M=\begin{bmatrix}1&0\\0&\boldsymbol{i}\end{bmatrix}$ *mod* 2Λ*. Then the characteristic polynomial of the endomorphism $\widetilde{f}^*_M$ of* $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ *is equal to*

$$\Big(\chi_{\det M^{-1}\times\mathrm{Sym}^2 M^{-1}}(x)\Big)^4\big(x-\det M\big)^3\big(x+\det M\big)^3.$$

In particular, if 1 *is not an eigenvalue of* $\det M\times\mathrm{Sym}^2 M$*, then $\widetilde{f}_M$ is infinitesimally rigid.*

4. Realization of infinitesimal deformations using divisors

Let X be a rational surface and let f be an automorphism of X. We will present a practical method to compute the actions f^* and f_* of f on $\mathrm{H}^1\big(X,\mathrm{T}X\big)$ using divisors. This is much more delicate than for the action on the Picard group. In practical examples, this method is effective (*see* §5).

4.1. 1-exceptional divisors.

— We start by defining a special class of divisors which will be of crucial importance in our study:

Definition 4.1. — Let X be a complex surface and let D be an effective divisor. We say that D is 1-*exceptional* if the natural morphism $\mathrm{H}^1\big(X,\mathrm{T}X\big)\longrightarrow \mathrm{H}^1\big(X,\mathrm{T}X(D)\big)$ vanishes.

The importance of these divisors comes from the following immediate observation: if D is 1-exceptional, then we have an exact sequence

$$0\longrightarrow \mathrm{H}^0\big(X,\mathrm{T}X\big)\longrightarrow \mathrm{H}^0\big(X,\mathrm{T}X(D)\big)\longrightarrow \mathrm{H}^0\big(D,\mathrm{T}X(D)_{|D}\big)\longrightarrow \mathrm{H}^1\big(X,\mathrm{T}X\big)\longrightarrow 0 \tag{4.1}$$

so that if X has no nonzero holomorphic vector field,

$$\mathrm{H}^1\big(X,\mathrm{T}X\big)\simeq \frac{\mathrm{H}^0\big(D,\mathrm{T}X(D)_{|D}\big)}{\mathrm{H}^0\big(X,\mathrm{T}X(D)\big)}.$$

Let us now explain how to construct these divisors on rational surfaces. We fix two complex surfaces X and Y such that Y is the blowup of X at a point p. If π is the blowup map and E is the exceptional divisor, we have a fundamental exact sequence relating $\mathrm{T}X$ and $\mathrm{T}Y$ via the differential of π which is

$$0\longrightarrow \mathrm{T}Y\longrightarrow \pi^*\mathrm{T}X\longrightarrow \iota_{E*}\Omega^1_E(-1)\longrightarrow 0. \tag{4.2}$$

For any locally free sheaf $\mathcal{F}$ on X, the natural pullback morphism $\mathrm{H}^i\big(X,\mathcal{F}\big)\longrightarrow \mathrm{H}^i\big(Y,\pi^*\mathcal{F}\big)$ is an isomorphism; this follows by writing down the Leray spectral sequence for the sheaf $\pi^*\mathcal{F}$ and using that $\mathrm{R}^j\pi_*\mathcal{O}_Y=0$ for $j>0$ (*see* Lemma 3.6 *(i)* where we already used this argument). Therefore, for every divisor D on X, we have a long exact sequence

$$0\longrightarrow \mathrm{H}^0\big(Y,\mathrm{T}Y(\pi^*D)\big)\longrightarrow \mathrm{H}^0\big(X,\mathrm{T}X(D)\big)\xrightarrow[\text{at}\,p]{\text{ev.}} \mathrm{T_pX}\otimes L_p\xrightarrow{\delta_D} \mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D)\big)\longrightarrow \mathrm{H}^1\big(X,\mathrm{T}X(D)\big)\longrightarrow 0$$

where δ_D is the connection morphism and L_p is the fiber of $\mathcal{O}_X(D)$ at p. We can now prove the first basic result about 1-exceptional divisors:

Proposition 4.2. — *If D is a* 1*-exceptional divisor on X, then π^*D+E is* 1*-exceptional on Y.*

Proof. — Let us start by proving that the composition of the connection morphism δ_D with the natural morphism $\mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D)\big)\longrightarrow \mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D+E)\big)$ vanishes. We have a morphism of short exact sequences

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{T}Y\big(\pi^*D\big) \arrow[r] \arrow[d] & \pi^*\mathrm{T}X\big(\pi^*D\big) \arrow[r] \arrow[d] & \iota_{E*}\Omega^1_E(-1)\otimes\mathcal{O}_Y\big(\pi^*D\big) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathrm{T}Y\big(\pi^*D+E\big) \arrow[r] & \pi^*\mathrm{T}X\big(\pi^*D+E\big) \arrow[r] & \iota_{E*}\Omega^1_E(-1)\otimes\mathcal{O}_Y\big(\pi^*D+E\big) \arrow[r] & 0
\end{tikzcd}$$

where the right vertical arrow vanishes. Therefore we get a commutative diagram

$$\begin{tikzcd}
\mathrm{H}^0\big(\iota_{E*}\Omega^1_E(-1)\otimes\mathcal{O}_Y(\pi^*D)\big) \arrow[r, "\delta_D"] \arrow[d] & \mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D)\big) \arrow[d] \\
\mathrm{H}^0\big(\iota_{E*}\Omega^1_E(-1)\otimes\mathcal{O}_Y(\pi^*D+E)\big) \arrow[r] & \mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D+E)\big)
\end{tikzcd}$$

and we get our claim since the left vertical arrow vanishes. We now consider the diagram

$$\begin{tikzcd}
 & \mathrm{H}^1\big(Y,\mathrm{T}Y\big) \arrow[r] \arrow[d, "u"] & \mathrm{H}^1\big(X,\mathrm{T}X\big) \arrow[d, "v"] \\
\mathrm{T}_pX\otimes L_p \arrow[r, "\delta_D"] & \mathrm{H}^1\big(Y,\mathrm{T}Y(\pi^*D)\big) \arrow[r] & \mathrm{H}^1\big(X,\mathrm{T}X(D)\big)
\end{tikzcd}$$

where the bottom line is exact. Since D is 1-exceptional, $v=0$. This implies that the image of u lies in the image of δ_D. □

Let us introduce some extra notations. For any effective divisor D on a surface X, we put:

$$\begin{cases} V(D) = \mathrm{H}^0\big(X, \mathrm{T}X(D)\big) \\ W(D) = \mathrm{H}^0\big(X, \mathrm{T}X(D)_{|D}\big). \end{cases}$$

We also denote by $\mathfrak{h}(\mathbb{P}^2)$ the set of holomorphic vector fields on $\mathbb{P}^2$.

Definition 4.3. — For any infinitely near point $\widehat{P}$ in $\mathbb{P}^2$ and any effective divisor D_{base} on $\mathbb{P}^2$, we define a divisor $D_{\widehat{P}, D_{\mathrm{base}}}$ on $\mathrm{Bl}_{\widehat{P}}\mathbb{P}^2$ as follows: if $\pi = \pi_N \circ \pi_{N-1} \circ \ldots \circ \pi_1$ is the sequence of blowups defining $\widehat{P}$ and $E_1, \ldots, E_N$ are the corresponding exceptional divisors, then $D_{\widehat{P}} = D_N$ where the D_i's are defined inductively by the formulas $D_0 = D_{\mathrm{base}}$ and $D_i = \pi_{i-1}^* D_{i-1} + E_i$ for $1 \leq i \leq N$. If D_{base} is empty, we put $D_{\varnothing, \widehat{P}} = D_{\widehat{P}}$.

According to Proposition 4.2 the divisor $D_{\widehat{P}, D_{\mathrm{base}}}$ is 1-exceptional on $\mathrm{Bl}_{\widehat{P}}\mathbb{P}^2$. Note that this definition extends readily to a finite number of infinitely near points in $\mathbb{P}^2$.

Lemma 4.4. — *For any infinitely near point $\widehat{P}$ in $\mathbb{P}^2$ of length N, the equality $\dim W(D_{\widehat{P}}) = 2N$ holds.*

Proof. — Let X be the rational surface obtained by blowing up $\widehat{P}$. Since $\pi(D_{\widehat{P}})$ is a finite set in $\mathbb{P}^2$, $V(D_{\widehat{P}})$ embeds in $\mathfrak{h}(\mathbb{P}^2)$. On the other hand, it is easy to see that each holomorphic vector field on $\mathbb{P}^2$ yields a meromorphic vector field on X which is in $V(D_{\widehat{P}})$. Therefore $V(D_{\widehat{P}})$ is isomorphic to $\mathfrak{h}(\mathbb{P}^2)$ so that we have

$$\mathrm{h}^1\big(X, \mathrm{T}X\big) = \dim W(D_{\widehat{P}}) - \Big(8 - \mathrm{h}^0\big(X, \mathrm{T}X\big)\Big).$$

Since $\mathrm{h}^1\big(X, \mathrm{T}X\big) - \mathrm{h}^0\big(X, \mathrm{T}X\big) = 2N - 8$, we get the result. $\square$

Proposition 4.5. — *Let X be a rational surface without nonzero holomorphic vector field obtained by blowing $\mathbb{P}^2$ in infinitely near points $\widehat{P}_1, \ldots, \widehat{P}_k$. Let D_{base} be an effective divisor on $\mathbb{P}^2$, and let $V^{\dagger}(D_{\mathrm{base}})$ be a direct factor of $\mathfrak{h}(\mathbb{P}^2)$ in $V(D_{\mathrm{base}})$. We denote by $\widehat{D}$ the 1-exceptional divisor $D_{\widehat{P}_1 \cup \ldots \cup \widehat{P}_k, D_{\mathrm{base}}}$. Then there is a natural isomorphism between $V(D_{\mathrm{base}})$ and $V(\widehat{D})$, and the associated morphism*

$$\bigoplus_{i=1}^{k} W(D_{\widehat{P}_i}) \oplus V^{\dagger}(D_{\mathrm{base}}) \longrightarrow W(\widehat{D})$$

is an isomorphism.

Proof. — For the first isomorphism, we argue exactly as in the proof of Lemma 4.4. For the second isomorphism, set $K = D_{\widehat{P}_1} + \ldots + D_{\widehat{P}_k}$. Let us write down the exact sequence (4.1) for the divisors K and $\widehat{D}$. Since $V(K) = \mathfrak{h}(\mathbb{P}^2)$, we get a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathfrak{h}(\mathbb{P}^2) \arrow[r] \arrow[d] & W(K) \arrow[r] \arrow[d] & \mathrm{H}^1\big(X, \mathrm{T}X\big) \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & V(D_{\mathrm{base}}) \arrow[r] & W(\widehat{D}) \arrow[r] & \mathrm{H}^1\big(X, \mathrm{T}X\big) \arrow[r] & 0
\end{tikzcd}$$

As $V(D_{\mathrm{base}}) = \mathfrak{h}(\mathbb{P}^2) \oplus V^{\dagger}(D_{\mathrm{base}})$, we obtain that $W(\widehat{D})$ is isomorphic to $W(K) \oplus V^{\dagger}(D_{\mathrm{base}})$. $\square$

We end the section with a statement which is particularly useful in practical computations:

Lemma 4.6. — *Let D, D' be two divisors on a rational surface X such that:*

$$\begin{cases} D' \textit{ is } 1-\textit{exceptional.} \\ \textit{The natural map from } W(D) \text{ to } W(D') \textit{ is surjective.} \end{cases}$$

Then D is 1-exceptional.

Proof. — We have a commutative diagram

$$\begin{tikzcd} W(D) \arrow[r] \arrow[d] & \mathrm{H}^1\big(X, \mathrm{T}X\big) \arrow[d, equal] \\ W(D') \arrow[r] & \mathrm{H}^1\big(X, \mathrm{T}X\big) \end{tikzcd}$$

where the bottom horizontal arrow is onto. Hence the top horizontal arrow is also onto. □

4.2. Geometric bases. — In this section, we construct a basis for the vector space $W(\widehat{D})$ using coverings (the divisor $\widehat{D}$ has been introduced in Proposition 4.5), we will call such a basis a *geometric basis*. Since $\mathrm{T}X(D)_{|D}$ is the sheaf quotient $\frac{\mathrm{T}X(D)}{\mathrm{T}X}$, a global section can be represented as a family of sections $(Z_i)_{i\in I}$ of $\mathrm{T}X(D)$ on open sets U_i covering D such that $Z_i - Z_j$ is holomorphic on U_{ij} for all i, j in I.

We start with the simplest case, namely $D_{\text{base}} = \varnothing$ and $k = 1$, so that $\widehat{D} = D_{\widehat{P}}$. Let us construct this basis by induction on the length N of $\widehat{P}$.

Let $\widehat{P}$ be an infinitely near point of $\mathbb{P}^2$ of length N, let q be a point in one of the exceptional divisors, $\widehat{P}' = \widehat{P} \cup \{q\}$, and put $X = \mathrm{Bl}_{\widehat{P}}\mathbb{P}^2$ and $Y = \mathrm{Bl}_{\widehat{P}'}\mathbb{P}^2$. Assume that we are given a covering of $D_{\widehat{P}}$ by open sets $(U_i)_{i\in I}$ of X such that :

① there exists a unique i_0 in I such that $q \in I_0$;

② the evaluation map $\mathrm{H}^0\big(U_{i_0}, \mathrm{T}U_{i_0}\big) \longrightarrow \mathrm{T}_qX$ is surjective;

③ there exists a basis $Z_1, \ldots, Z_{2N}$ of $W(D_{\widehat{P}})$, where for each α, Z_α is a collection of holomorphic sections $(Z_{i\alpha})_{i\in I}$ of $\mathrm{T}X(D_{\widehat{P}})$ on the U_i's such that for all i and j, the section $Z_{i\alpha} - Z_{j\alpha}$ is holomorphic on each $U_{i\alpha} \cap U_{j\alpha}$.

Let $\pi\colon Y \to X$ be the blowup map and let E be the exceptional divisor. Set $U'_i = \pi^{-1}(U_i)$. For any basis (v_1, v_2) of $\mathrm{T}_q\left(\mathrm{Bl}_{\widehat{P}}\mathbb{P}^2\right)$, thanks to ②, we can choose two holomorphic vector fields T_1, T_2 on U_{i_0} which extend (v_1, v_2). We will consider the vector fields T_1 and T_2 as sections of $\mathrm{T}Y(E)$ on U'_{i_0}. Let us now consider the sections $(Z'_1, \ldots, Z'_{2N+2})$ of $W\big(D_{\widehat{P}'}\big)$ defined using the covering $(U'_i)_{i\in I}$ as follows :

	Z'_1	……	Z'_{2N}	Z'_{2N+1}	Z'_{2N+2}
U'_{i_0}	Z_{1i_0}	……	Z_{2Ni_0}	T_1	T_2
$U'_j \quad j \neq i_0$	Z_{1j}	……	Z_{2Nj}	0	0

Then we have the following result:

Proposition 4.7. — *The family $(Z'_1, \ldots, Z'_{2N+2})$ is a basis of $W\big(D_{\widehat{P}'}\big)$.*

Proof. — Thanks to Lemma 4.4, it suffices to show that this family is free. Let $\lambda_1, \ldots, \lambda_{2N+2}$ be complex numbers such that $\lambda_1 Z'_1 + \lambda_2 Z'_2 + \ldots + \lambda_{2N+2} Z'_{2N+2} = 0$. Then for all $j \neq i_0$,

$$\lambda_1 Z'_{1j} + \lambda_2 Z'_{2j} + \ldots + \lambda_{2N} Z'_{2Nj}$$

is holomorphic on U'_j, and $\lambda_1 Z'_{1i_0} + \lambda_2 Z'_{2i_0} + \ldots + \lambda_{2N} Z'_{2Ni_0} + \lambda_{2N+1} T_1 + \lambda_{2N+2} T_2$ is holomorphic on U'_{i_0}.

Therefore, $\lambda_1 Z'_{1j} + \lambda_2 Z'_{2j} + \ldots + \lambda_{2N} Z'_{2Nj}$ is holomorphic on U'_j also for $j = i_0$ and $\lambda_{2N+1} T_1 + \lambda_{2N+2} T_2$ (considered as a vector field on U_{i_0}) vanishes at q. Since $(Z_1, \ldots, Z_{2N})$ is a basis of $W(D_{\widehat{P}})$ we get $\lambda_1 = \ldots = \lambda_{2N} = 0$; and as (v_1, v_2) is a basis of T_qX, $\lambda_{2N+1} = \lambda_{2N+2} = 0$. □

Remarks 4.8. —

(i) In practical computations, U_{i_0} is a domain for some holomorphic coordinates (z, w), $T_1 = \partial_z$ and $T_2 = \partial_w$.

(ii) If q' is a point on $\widehat{P}'$, the covering $(U_i')_{i\in I}$ does not satisfy in general conditions ② and ③. Therefore it is necessary to refine the covering at each blowup.

Using Proposition 4.5, we can now easily construct a basis of $W\big(\widehat{D}\big)$. Indeed, we have already explained how to construct a basis of each $W\big(D_{\widehat{P}_k}\big)$. Then it suffices to take a basis of $V(D_{\text{base}})^{\dagger}$ and to pull back these meromorphic vector fields to the surface.

4.3. Algebraic bases. — In practical computations, if an element of $W(\widehat{D})$ is given by local meromorphic vector fields on open sets of a covering, it is not a priori obvious to decompose this element in a geometric basis (as constructed in §4.2). In this section, we will construct another basis (we call it an algebraic basis) which solves this problem.

We start by defining the residue morphisms. Let $\widehat{P}=\{p_1,\dots,p_N\}$ and let $\overline{E}_1,\dots,\overline{E}_N$ be the corresponding strict transforms of the exceptional divisors. We fix holomorphic coordinates (x_i,y_i) near p_i such that $p_i=(0,0)$ in these coordinates, and we introduce new holomorphic coordinates (u_i,v_i) by putting $x_i=u_i$ and $y_i=u_iv_i$. We can consider (u_i,v_i) as holomorphic coordinates on an open subset of X which contains $\overline{E}_i$ except a finite number of points.

Let $(0,\lambda_i)$ be a generic point of $\overline{E}_i$, let Z be an element of $W\big(D_{\widehat{P}}\big)$ and let Z_i be a section of $\mathrm{T}X(D)$ which lifts Z near this point. Then Z_i admits a Laurent expansion

$$Z_i(u_i,v_i)=\sum_{n=1}^{n_0}\sum_{m=0}^{\infty}\frac{(v_i-\lambda_i)^m(a_{nm}\,\partial_u+b_{nm}\,\partial_v)}{u^n}+\text{holomorphic terms.}$$

Definition 4.9. — For any generic complex number λ_i, we define the i^{th} *residue morphism* $\mathrm{res}_{E_i}\colon W(D_{\widehat{P}})\longrightarrow\mathbb{C}^2$ by the formula $\mathrm{res}_{E_i}(Z)=(b_{10},b_{11})$. We also define $\mathrm{res}_{\widehat{P}}\colon W(D_{\widehat{P}})\longrightarrow\mathbb{C}^{2N}$ by $\mathrm{res}_{\widehat{P}}=\bigoplus_{i=1}^{N}\mathrm{res}_{E_i}$.

Note that the definition of these residues depends on the coordinates and on the generic parameters on the exceptional divisors.

Lemma 4.10. — *The morphism* $\mathrm{res}_{\widehat{P}}$ *is an isomorphism.*

Proof. — Let us take a geometric basis associated to the coordinates $(x_i,y_i)_{1\leq i\leq N}$ as constructed in §4.2. Then the matrix of $\mathrm{res}_{\widehat{P}}$ is lower triangular by blocks, where the diagonal blocks are the 2×2 matrices whose columns are the vectors $\mathrm{res}_{E_i}(\partial_{x_i})$ and $\mathrm{res}_{E_i}(\partial_{y_i})$. We have $\partial_{x_i}=\partial_{u_i}-\frac{v_i}{u_i}\partial_{v_i}$ and $\partial_{y_i}=\frac{1}{u_i}\partial_{v_i}$. Therefore we obtain $\mathrm{res}_{E_i}(\partial_{x_i})=(-1,-\lambda_i)$ and $\mathrm{res}_{E_i}(\partial_{y_i})=(1,0)$, so that the diagonal blocks are all invertible. ☐

We can now define algebraic basis in the general case. We take the notation of Proposition 4.5. In order to avoid cumbersome notation, we will assume for simplicity that D_{base} is irreducible, and leave the general case to the reader.

Let $\chi_1,\dots,\chi_r$ be a basis of $V(D_{\text{base}})^{\dagger}$.

Lemma 4.11. — *Assume that* D_{base} *is irreducible, and let* Z *be an element of* $W(\widehat{D})$. *Then there exist unique complex numbers* $\alpha_1(Z),\dots,\alpha_r(Z)$ *such that for any generic point* ξ *of* D_{base} *and any lift* $\widetilde{Z}$ *of* Z *near* ξ,

$$\widetilde{Z}=\alpha_1(Z)\,\chi_1+\ldots+\alpha_m(Z)\,\chi_r+\text{ holomorphic terms}$$

in a neighborhood of ξ.

Proof. — The existence of such a decomposition is straightforward: the $\alpha_i(Z)$'s are the coefficients of Z in $V(D_{\text{base}})^{\dagger}$ when decomposed in a geometric basis. For the unicity, we remark that $\alpha_1\,\chi_1+\ldots+\alpha_r\,\chi_r$ is holomorphic near a point of D_{base}, it must be holomorphic on $\mathbb{P}^2$ since D_{base} is irreducible. Since $V(D_{\text{base}})^{\dagger}$ is a direct factor of $\mathfrak{h}(\mathbb{P}^2)$, all the coefficients α_i must vanish. ☐

Corollary 4.12. — *With the notations of Proposition 4.5, the map*

$$\big(\mathrm{res}_{\widehat{P}_1},\dots,\mathrm{res}_{\widehat{P}_k},\alpha_1,\dots,\alpha_r\big)\colon W(\widehat{D})\longrightarrow\mathbb{C}^{2N_1+\ldots+2N_k+r}$$

is an isomorphism.

By definition, the associated algebraic basis of $W(\widehat{D})$ is the image of the canonical basis of $\mathbb{C}^{2N_1+\ldots+2N_k+r}$ by the inverse isomorphism.

5. An explicit example on $\mathbb{P}^2$ blown up in 15 points

5.1. The strategy. — Let $\phi\colon \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ be the birational map given by $[x\colon y\colon z] \to [xz^2+y^3\colon yz^2\colon z^3]$. The map ϕ induces an isomorphism (which we still denote by ϕ) between $\mathrm{Bl}_{\widehat{P}_1}\,\mathbb{P}^2$ and $\mathrm{Bl}_{\widehat{P}_2}\,\mathbb{P}^2$, where $\widehat{P}_1$ and $\widehat{P}_2$ are infinitely near points of $\mathbb{P}^2$ of length 5 centered at $[1:0:0]$ described in [**7**, §2.1]. For any complex number α, we put

$$A = \begin{bmatrix} \alpha & 2(1-\alpha) & 2+\alpha-\alpha^2 \\ -1 & 0 & \alpha+1 \\ 1 & -2 & 1-\alpha \end{bmatrix}$$

and we consider it as an element of $\mathrm{PGL}(3;\mathbb{C})$. Then the map $A\phi$ lifts to an automorphism ψ of the rational surface X obtained by blowing up the projective plane 15 times at $\widehat{P}_1$, $A\widehat{P}_2$ and $A\phi A\widehat{P}_2$ (*see* [**7**, §3.3]). Recall that the parameter α is not really interesting because two different values of α correspond to linearly conjugate automorphisms, however we keep it in order to check some calculations in the sequel. Our aim is to compute the map ψ_* acting on $\mathrm{H}^1(X,\mathrm{T}X)$.

Let us recall the construction of $\widehat{P}_1$ $\big($resp. $\widehat{P}_2\big)$. In affine coordinates (y,z), we blow up the point $(0,0)_{y,z}$ in $\mathbb{P}^2$ and we denote by E the exceptional divisor. We put $y=u_1$, $z=u_1v_1$, we blow up $(0,0)_{u_1,v_1}$ and F is the exceptional divisor. Then we put $u_1=r_2s_2$, $v_1=s_2$, we blow up $(0,0)_{r_2,s_2}$ and we denote by G the exceptional divisor. Next, we put $r_2=r_3s_3$, $s_2=s_3$, we blow up $(-1,0)_{r_3,s_3}$ $\big($resp. $(1,0)_{r_3,s_3}\big)$ and H (resp. K) is the exceptional divisor. Then, we put $r_3=r_4s_4-1$, $s_3=s_4$ $\big($resp. $r_3=c_4d_4+1$, $s_3=d_4\big)$, we blow up $(0,0)_{r_4,s_4}$ $\big($resp. $(0,0)_{c_4,d_4}\big)$ and we denote by L (resp. M) the last exceptional divisor. Lastly, we put $r_4=r_5s_5$, $s_4=s_5$ $\big($resp. $c_4=c_5d_5$, $d_4=d_5\big)$.

Set $X_1=\mathrm{Bl}_{\widehat{P}_1}\,\mathbb{P}^2$ and $X_2=\mathrm{Bl}_{\widehat{P}_2}\,\mathbb{P}^2$. Let $\overline{\Delta}$ be the strict transform of the line $\Delta=\{z=0\}$ in X_2. Then

$$\phi(E)=E,\quad \phi(F)=K,\quad \phi(G)=G,\quad \phi(H)=F,\quad \phi(L)=\overline{\Delta}. \tag{5.1}$$

We have $D_{\widehat{P}_1}=E+2F+4G+5H+6L$ and $D_{\widehat{P}_2}=E+2F+4G+5K+6M$. Let D_1 be the divisor on X_1 given by

$$D_1=E+2F+3G+4H+5L.$$

By explicit calculation, the natural morphism from $W(D_1)$ to $W\big(D_{\widehat{P}_1}\big)$ is surjective so that D_1 is 1-exceptional on X_1 by Lemma 4.6. We now define a divisor D_2 on X_2 as follows (*see* Definition 4.3):

$$D_2=D_{\widehat{P}_2,5\overline{\Delta}}.$$

By (5.1), we have $\phi_*D_1\leq D_2$. Let $\mathfrak{D}_1$ and $\mathfrak{D}_2$ be the two 1-exceptional divisors on X given by:

$$\mathfrak{D}_1=D_1+AD_{\widehat{P}_2}+A\phi A\,D_{\widehat{P}_2},\qquad\qquad \mathfrak{D}_2=D_{\widehat{P}_1}+AD_2+A\phi A\,D_{\widehat{P}_2}.$$

Then $\mathfrak{D}_1\leq\mathfrak{D}_2$ and $f_*\mathfrak{D}_1\leq\mathfrak{D}_2$. Therefore the morphism ψ_* acting on $\mathrm{H}^1(X,\mathrm{T}X)$ can be obtained as the composition

$$\frac{W(\mathfrak{D}_1)}{V(\mathfrak{D}_1)}\xrightarrow{\ \psi_*\ }\frac{W(\mathfrak{D}_2)}{V(\mathfrak{D}_2)}\xrightarrow{\ \sim\ }\frac{W(\mathfrak{D}_1)}{V(\mathfrak{D}_1)} \tag{5.2}$$

where the inverse of last arrow is induced by the natural morphism from $W(\mathfrak{D}_1)$ to $W(\mathfrak{D}_2)$. To compute ψ_*, the strategy runs as follows:

Calculations for the pair (D_1,D_2).

Step 1 – Express the vectors of a geometric basis of $W(D_1)$ in an algebraic basis.

Step 2 – Compute $\phi_*\colon W(D_1)\longrightarrow W(D_2)$ where $W(D_1)$ is endowed with a geometric basis and $W(D_2)$ is endowed with an algebraic basis.

Step 3 – Find bases of $V(D_1)$ and $V(D_2)$ whose vectors are expressed in algebraic bases of $W(D_1)$ and $W(D_2)$ respectively.

Calculations for the pair $(\mathfrak{D}_1,\mathfrak{D}_2)$.

Step 1 – Compute $\psi_*\colon W(\mathfrak{D}_1)\longrightarrow W(\mathfrak{D}_2)$ where $W(\mathfrak{D}_1)$ and $W(\mathfrak{D}_2)$ are endowed with algebraic bases.
Step 2 – Find bases of $V(\mathfrak{D}_1)$ and $V(\mathfrak{D}_2)$ whose vectors are expressed in algebraic bases of $W(\mathfrak{D}_1)$ and $W(\mathfrak{D}_2)$ respectively.

5.2. Calculations for the pair (D_1,D_2). — We give results of the calculations for the first three steps listed above.

Step 1 – We use the coordinates (y,z), (u_1,v_1), (r_2,s_2), (r_3,s_3), (r_4,s_4) and (r_5,s_5) to compute the algebraic and geometric bases on X_1. Let $(U_i)_{1\leq i\leq 5}$ be the covering of D_1 given by the following picture:



Then a geometric basis of $W(D_1)$ is given by the ten following vectors:

	U_1	U_2	U_3	U_4	U_5
$\mathfrak{e}_1$	∂y	∂y	∂y	∂y	∂y
$\mathfrak{e}_2$	∂z	∂z	∂z	∂z	∂z
$\mathfrak{e}_3$	0	∂u_1	∂u_1	∂u_1	∂u_1
$\mathfrak{e}_4$	0	∂v_1	∂v_1	∂v_1	∂v_1
$\mathfrak{e}_5$	0	0	∂r_2	∂r_2	∂r_2
$\mathfrak{e}_6$	0	0	∂s_2	∂s_2	∂s_2
$\mathfrak{e}_7$	0	0	0	∂r_3	∂r_3
$\mathfrak{e}_8$	0	0	0	∂s_3	∂s_3
$\mathfrak{e}_9$	0	0	0	0	∂r_4
$\mathfrak{e}_{10}$	0	0	0	0	∂s_4

– Matrix $\mathscr{P}$ –

If λ_1, λ_2, λ_3, λ_4 and λ_5 are generic parameters on E, F, G, H and L respectively and if $(\mathfrak{l}_i)_{1\leq i\leq 10}$ is the associated algebraic basis, a direct calculation yields:

	$\mathfrak{e}_1$	$\mathfrak{e}_2$	$\mathfrak{e}_3$	$\mathfrak{e}_4$	$\mathfrak{e}_5$	$\mathfrak{e}_6$	$\mathfrak{e}_7$	$\mathfrak{e}_8$	$\mathfrak{e}_9$	$\mathfrak{e}_{10}$
$\mathfrak{l}_1$	$-\lambda_1$	1	0	0	0	0	0	0	0	0
$\mathfrak{l}_2$	-1	0	0	0	0	0	0	0	0	0
$\mathfrak{l}_3$	2	0	1	$-\lambda_2$	0	0	0	0	0	0
$\mathfrak{l}_4$	0	0	0	-1	0	0	0	0	0	0
$\mathfrak{l}_5$	0	0	0	$-2\lambda_3$	1	$-\lambda_3$	0	0	0	0
$\mathfrak{l}_6$	0	0	0	-2	0	-1	0	0	0	0
$\mathfrak{l}_7$	$-\lambda_4^2$	λ_4^3	0	$-3\lambda_4$	0	$-2\lambda_4$	1	$-\lambda_4$	0	0
$\mathfrak{l}_8$	$-2\lambda_4$	$3\lambda_4^2$	0	-3	0	-2	0	-1	0	0
$\mathfrak{l}_9$	0	$2\lambda_5^2$	0	$-4\lambda_5$	0	$-3\lambda_5$	0	$-2\lambda_5$	1	$-\lambda_5$
$\mathfrak{l}_{10}$	0	$4\lambda_5$	0	-4	0	-3	0	-2	0	-1

– *Matrix* $\mathscr{K}$ –

Step 2 – We start by fixing a basis of $\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2(m\Delta)\big)$. We divide $\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2(m\Delta)\big)$ in four subspaces, the first one corresponding to holomorphic vector fields.

–Vector fields of type a that span a subspace of dimension 8:

$$a_1=\partial_y,\quad a_2=y\partial_y,\quad a_3=z\partial_y,\quad a_4=\partial_z,\quad a_5=y\partial_z,\quad a_6=z\partial_z,\quad a_7=yz\partial_z,\quad a_8=z^2\partial_z.$$

–Vector fields of type b that span a subspace of dimension $\frac{m(m+5)}{2}$:

$$b_{p,q}=\frac{y^p}{z^q}\partial_y \qquad 1\leq q\leq m,\quad 0\leq p\leq q+1.$$

–Vector fields of type c that span a subspace of dimension $\frac{m(m+5)}{2}$:

$$c_{p,q}=\frac{y^p}{z^q}\partial_z \qquad 1\leq q\leq m,\quad 0\leq p\leq q+1.$$

–Vector fields of type d that span a subspace of dimension m:

$$d_p=\frac{y^{p+2}}{z^p}\partial_y+\frac{y^{p+1}}{z^{p-1}}\partial_z \qquad 1\leq p\leq m.$$

We take for $\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2(m\Delta)\big)^\dagger$ the subspace spanned by meromorphic vector fields of type b, c, and d (in our setting, $m=5$). Then we use the coordinates (y,z), (u_1,v_1), (r_2,s_2), (r_3,s_3), (c_4,d_4) and (c_5,d_5) on X_2. If μ_1, μ_2, μ_3, μ_4 and μ_5 are the generic parameters on E, F, G, K and M respectively, we consider an algebraic basis of $W(D_2)$ associated with these parameters consisting of ten vectors $(\mathfrak{m}_i)_{1\leq i\leq 10}$ corresponding to the exceptional divisors, and 55 vectors (25 of type b, 25 of type c and 5 of type d) corresponding to a basis of $\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2(5\Delta)\big)^\dagger$.

By direct computation, we have

$$\phi_*\partial_y=\left(1-\frac{4y^3}{z^2}+\frac{3y^6}{z^4}\right)\partial_y+\left(-\frac{3y^2}{z}+\frac{3y^5}{z^3}\right)\partial_z, \qquad \phi_*\partial_z=2\left(\frac{y^4}{z^3}-\frac{y^7}{z^5}\right)\partial_y+\left(1+\frac{y^3}{z^2}-\frac{2y^6}{z^4}\right)\partial_z,$$

$$\phi_*\partial_{u_1}=\left(1-2\frac{y^3}{z^2}+\frac{y^6}{z^4}\right)\partial_y+\left(\frac{z}{y}-2\frac{y^2}{z}+\frac{y^5}{z^3}\right)\partial_z, \qquad \phi_*\partial_{v_1}=2\frac{y^5}{z^3}\partial_y+\left(y+2\frac{y^4}{z^2}\right)\partial_z,$$

$$\phi_*\partial_{r_2}=\left(\frac{z}{y}-2\frac{y^2}{z}+\frac{y^5}{z^3}\right)\partial_y+\left(\frac{z^2}{y^2}-2y+\frac{y^4}{z^2}\right)\partial_z, \qquad \phi_*\partial_{s_2}=\left(\frac{y^2}{z}+\frac{y^5}{z^3}\right)\partial_y+\left(2y+\frac{y^4}{z^2}\right)\partial_z,$$

$$\phi_*\partial_{r_3}=\left(\frac{z^2}{y^2}-2y+\frac{y^4}{z^2}\right)\partial_y+\left(\frac{z^3}{y^3}-2z+\frac{y^3}{z}\right)\partial_z, \qquad \phi_*\partial_{s_3}=\frac{2y^2}{z}\partial_y+3y\partial_z,$$

$$\phi_*\partial_{r_4}=\left(\frac{z^3}{y^3}-2z+\frac{y^3}{z}\right)\partial_y+\left(\frac{z^4}{y^4}-\frac{2z^2}{y}+y^2\right)\partial_z, \qquad \phi_*\partial_{s_4}=\left(\frac{z}{y}+\frac{y^2}{z}\right)\partial_y+\left(\frac{z^2}{y^2}+2y\right)\partial_z.$$

Therefore, the matrix of $\phi_* \colon W(D_1) \longrightarrow W(D_2)$ is:

	$\phi_*\mathfrak{e}_1$	$\phi_*\mathfrak{e}_2$	$\phi_*\mathfrak{e}_3$	$\phi_*\mathfrak{e}_4$	$\phi_*\mathfrak{e}_5$	$\phi_*\mathfrak{e}_6$	$\phi_*\mathfrak{e}_7$	$\phi_*\mathfrak{e}_8$	$\phi_*\mathfrak{e}_9$	$\phi_*\mathfrak{e}_{10}$
$\mathfrak{m}_1$	$-\mu_1$	1	0	0	0	0	0	0	0	0
$\mathfrak{m}_2$	-1	0	0	0	0	0		0	0	0
$\mathfrak{m}_3$	2	0	1	$-\mu_2$	$-2\mu_2$	0	μ_2^2	μ_2	0	0
$\mathfrak{m}_4$	0	0	0	-1	-2	0	$2\mu_2$	1	0	0
$\mathfrak{m}_5$	0	0	0	$2\mu_3(\mu_3-1)$	$(\mu_3-1)^2$	$\mu_3(\mu_3-1)$	0	0	0	0
$\mathfrak{m}_6$	0	0	0	$4\mu_3-2$	$2(\mu_3-1)$	$2\mu_3-1$	0	0	0	0
$\mathfrak{m}_7$	$2\mu_4^2$	$-\mu_4^3$	μ_4^2	μ_4	0	0	0	0	0	0
$\mathfrak{m}_8$	$4\mu_4$	$-3\mu_4^2$	$2\mu_4$	1	0	0	0	0	0	0
$\mathfrak{m}_9$	0	0	0	0	0	0	0	0	0	0
$\mathfrak{m}_{10}$	0	0	0	0	0	0	0	0	0	0
$\mathrm{H}^0\big(\mathbb{P}^2, \mathrm{T}\mathbb{P}^2(5\Delta)\big)^\dagger$	$-4b_{3,2}$ $-3c_{2,1}$ $3d_4$	$2b_{4,3}$ $c_{3,2}$ $-2d_5$	$-2b_{3,2}$ $-2c_{2,1}$ d_4	$2d_3$	$-2b_{2,1}$ d_3	$b_{2,1}$ d_3	d_2	$2b_{2,1}$	d_1	$b_{2,1}$

– Matrix $\mathscr{L}$ –

Step 3 – For typographical reasons, we will write the vectors of $V(D_1)$ and $V(D_2)$ as lines (and not as columns).

	$\mathfrak{l}_1$	$\mathfrak{l}_2$	$\mathfrak{l}_3$	$\mathfrak{l}_4$	$\mathfrak{l}_5$	$\mathfrak{l}_6$	$\mathfrak{l}_7$	$\mathfrak{l}_8$	$\mathfrak{l}_9$	$\mathfrak{l}_{10}$
a_1	$-\lambda_1$	-1	2	0	0	0	$-\lambda_4^2$	$-2\lambda_4$	0	0
a_2	0	0	0	0	0	0	-3	0	0	0
a_3	0	0	0	0	0	0	0	0	-3	0
a_4	1	0	0	0	0	0	λ_4^3	$3\lambda_4^2$	$2\lambda_5^2$	$4\lambda_5$
a_5	0	0	$-\lambda_2$	-1	$-2\lambda_3$	-2	$-3\lambda_4$	-3	$-4\lambda_5$	-4
a_6	0	0	0	0	0	0	2	0	0	0
a_7	0	0	0	0	0	0	0	0	0	0
a_8	0	0	0	0	0	0	0	0	0	0

– Matrix ${}^t\mathscr{M}$ –

	$\mathfrak{m}_1$	$\mathfrak{m}_2$	$\mathfrak{m}_3$	$\mathfrak{m}_4$	$\mathfrak{m}_5$	$\mathfrak{m}_6$	$\mathfrak{m}_7$	$\mathfrak{m}_8$	$\mathfrak{m}_9$	$\mathfrak{m}_{10}$
a_1	$-\mu_1$	-1	2	0	0	0	$-\mu_4^2$	$-2\mu_4$	0	0
a_2	0	0	0	0	0	0	3	0	0	0
a_3	0	0	0	0	0	0	0	0	3	0
a_4	1	0	0	0	0	0	$-\mu_4^3$	$-3\mu_4^2$	$2\mu_5^2$	$4\mu_5$
a_5	0	0	$-\mu_2$	-1	$-2\mu_3$	-2	$-3\mu_4$	-3	$-4\mu_5$	-4
a_6	0	0	0	0	0	0	-2	0	0	0
a_7	0	0	0	0	0	0	0	0	0	0
a_8	0	0	0	0	0	0	0	0	0	0

– Matrix ${}^t\mathscr{N}_{\mathrm{a}}$ –

	$\mathfrak{m}_1$	$\mathfrak{m}_2$	$\mathfrak{m}_3$	$\mathfrak{m}_4$	$\mathfrak{m}_5$	$\mathfrak{m}_6$	$\mathfrak{m}_7$	$\mathfrak{m}_8$	$\mathfrak{m}_9$	$\mathfrak{m}_{10}$
$b_{0,1}$	0	0	0	0	0	0	$2\mu_4^5$	$10\mu_4^4$	$3\mu_5^3$	$9\mu_5^2$
$b_{1,1}$	-1	0	0	0	0	0	μ_4^3	$3\mu_4^2$	$-2\mu_5^2$	$-4\mu_5$
$b_{2,1}$	0	0	$2\mu_2$	2	$3\mu_3$	3	$4\mu_4$	4	$5\mu_5$	5
$b_{0,2}$	0	0	0	0	0	0	$-9\mu_4^8$	$-72\mu_4^7$	0	0
$b_{1,2}$	0	0	0	0	0	0	$-3\mu_4^6$	$-18\mu_4^5$	0	0
$b_{2,2}$	$-\mu_1{}^{-1}$	μ_1^{-2}	0	0	0	0	$-\mu_4^4$	$-4\mu_4^3$	0	0
$b_{3,2}$	0	0	0	0	0	0	0	0	0	0
$b_{0,3}$	0	0	0	0	0	0	$52\mu_4^{11}$	$572\mu_4^{10}$	$-28\mu_5^6$	$-168\mu_5^5$
$b_{1,3}$	0	0	0	0	0	0	$15\mu_4^9$	$135\mu_4^8$	$12\mu_5^5$	$60\mu_5^4$
$b_{2,3}$	0	0	0	0	0	0	$4\mu_4^7$	$28\mu_4^6$	$-5\mu_5^4$	$-20\mu_5^3$
$b_{3,3}$	$-\mu_1^{-2}$	$2\mu_1^{-3}$	0	0	0	0	μ_4^5	$5\mu_4^4$	$2\mu_5^3$	$6\mu_5^2$
$b_{4,3}$	0	0	0	0	0	0	0	0	0	0
$b_{0,4}$	0	0	0	0	0	0	$-340\mu_4^{14}$	$-4760\mu_4^{13}$	0	0
$b_{1,4}$	0	0	0	0	0	0	$-91\mu_4^{12}$	$-1092\mu_4^{11}$	0	0
$b_{2,4}$	0	0	0	0	0	0	$-22\mu_4^{10}$	$-220\mu_4^9$	0	0
$b_{3,4}$	0	0	0	0	0	0	$-5\mu_4^8$	$-40\mu_4^7$	0	0
$b_{4,4}$	$-\mu_1^{-3}$	$3\mu_1^{-4}$	0	0	0	0	$-\mu_4^6$	$-6\mu_4^5$	0	0
$b_{5,4}$	0	0	0	0	0	0	0	0	0	0
$b_{0,5}$	0	0	0	0	0	0	$2394\mu_4^{17}$	$40698\mu_4^{16}$	$429\mu_5^9$	$3861\mu_5^8$
$b_{1,5}$	0	0	0	0	0	0	$612\mu_4^{15}$	$9180\mu_4^{14}$	$-165\mu_5^8$	$-1320\mu_5^7$
$b_{2,5}$	0	0	0	0	0	0	$140\mu_4^{13}$	$1820\mu_4^{12}$	$60\mu_5^7$	$420\mu_5^6$
$b_{3,5}$	0	0	0	0	0	0	$30\mu_4^{11}$	$330\mu_4^{10}$	$-21\mu_5^6$	$-126\mu_5^5$
$b_{4,5}$	0	0	0	0	0	0	$6\mu_4^9$	$54\mu_4^8$	$7\mu_5^5$	$35\mu_5^4$
$b_{5,5}$	$-\mu_1^{-4}$	$4\mu_1^{-5}$	0	0	0	0	μ_4^7	$7\mu_4^6$	$-2\mu_5^4$	$-8\mu_5^3$
$b_{6,5}$	0	0	0	0	0	0	0	0	0	0

– *Matrix* ${}^t\mathscr{N}_{\mathfrak{b}}$ –

	$\mathfrak{m}_1$	$\mathfrak{m}_2$	$\mathfrak{m}_3$	$\mathfrak{m}_4$	$\mathfrak{m}_5$	$\mathfrak{m}_6$	$\mathfrak{m}_7$	$\mathfrak{m}_8$	$\mathfrak{m}_9$	$\mathfrak{m}_{10}$
$c_{0,1}$	0	0	0	0	0	0	$4\mu_4^6$	$24\mu_4^5$	0	0
$c_{1,1}$	μ_1^{-1}	$-\mu_1^{-2}$	0	0	0	0	μ_4^4	$4\mu_4^3$	0	0
$c_{2,1}$	0	0	0	0	0	0	0	0	0	0
$c_{0,2}$	0	0	0	0	0	0	$-25\mu_4^9$	$-225\mu_4^8$	$-18\mu_5^5$	$-90\mu_5^4$
$c_{1,2}$	0	0	0	0	0	0	$-5\mu_4^7$	$-35\mu_4^6$	$6\mu_5^4$	$24\mu_5^3$
$c_{2,2}$	μ_1^{-2}	$-2\mu_1^{-3}$	0	0	0	0	$-\mu_4^5$	$-5\mu_4^4$	$-2\mu_5^3$	$-6\mu_5^2$
$c_{3,2}$	0	0	0	0	0	0	0	0	0	0
$c_{0,3}$	0	0	0	0	0	0	$182\mu_4^{12}$	$2184\mu_4^{11}$	0	0
$c_{1,3}$	0	0	0	0	0	0	$33\mu_4^{10}$	$330\mu_4^9$	0	0
$c_{2,3}$	0	0	0	0	0	0	$6\mu_4^8$	$48\mu_4^7$	0	0
$c_{3,3}$	μ_1^{-3}	$-3\mu_1^{-4}$	0	0	0	0	μ_4^6	$6\mu_4^5$	0	0
$c_{4,3}$	0	0	0	0	0	0	0	0	0	0
$c_{0,4}$	0	0	0	0	0	0	$-1428\mu_4^{15}$	$-21420\mu_4^{14}$	$330\mu_5^8$	$2640\mu_5^7$
$c_{1,4}$	0	0	0	0	0	0	$-245\mu_4^{13}$	$-3185\mu_4^{12}$	$-96\mu_5^7$	$-672\mu_5^6$
$c_{2,4}$	0	0	0	0	0	0	$-42\mu_4^{11}$	$-462\mu_4^{10}$	$28\mu_5^6$	$168\mu_5^5$
$c_{3,4}$	0	0	0	0	0	0	$-7\mu_4^9$	$-63\mu_4^8$	$-8\mu_5^5$	$-40\mu_5^4$
$c_{4,4}$	μ_1^{-4}	$-4\mu_1^{-5}$	0	0	0	0	$-\mu_4^7$	$-7\mu_4^6$	$2\mu_5^4$	$8\mu_5^3$
$c_{5,4}$	0	0	0	0	0	0	0	0	0	0
$c_{0,5}$	0	0	0	0	0	0	$11704\mu_4^{18}$	$210672\mu_4^{17}$	0	0
$c_{1,5}$	0	0	0	0	0	0	$1938\mu_4^{16}$	$31008\mu_4^{15}$	0	0
$c_{2,5}$	0	0	0	0	0	0	$320\mu_4^{14}$	$4480\mu_4^{13}$	0	0
$c_{3,5}$	0	0	0	0	0	0	$52\mu_4^{12}$	$624\mu_4^{11}$	0	0
$c_{4,5}$	0	0	0	0	0	0	$8\mu_4^{10}$	$80\mu_4^9$	0	0
$c_{5,5}$	μ_1^{-5}	$-5\mu_1^{-6}$	0	0	0	0	μ_4^8	$8\mu_4^7$	0	0
$c_{6,5}$	0	0	0	0	0	0	0	0	0	0

– *Matrix* ${}^t\mathscr{N}_{\mathrm{c}}$ –

	$\mathfrak{m}_1$	$\mathfrak{m}_2$	$\mathfrak{m}_3$	$\mathfrak{m}_4$	$\mathfrak{m}_5$	$\mathfrak{m}_6$	$\mathfrak{m}_7$	$\mathfrak{m}_8$	$\mathfrak{m}_9$	$\mathfrak{m}_{10}$
d_1	0	0	0	0	0	0	0	0	1	0
d_2	0	0	μ_2^2	$2\mu_2$	0	0	1	0	0	0
d_3	0	0	0	0	μ_3^2	$2\mu_3$	$2\mu_4$	2	$2\mu_5$	2
d_4	0	0	0	0	0	0	μ_4^2	$2\mu_4$	0	0
d_5	0	0	0	0	0	0	0	0	μ_5^2	$2\mu_5$

– *Matrix* ${}^t\mathscr{N}_{\mathrm{d}}$ –

Although we won't need it, it is easy to verify that

$$\left\{\begin{array}{lll} \phi_*(a_1)=a_1-4b_{3,2}-3c_{2,1}+3d_4, & \phi_*(a_2)=a_2-3d_2, & \phi_*(a_3)=a_3-3d_1,\\ \phi_*(a_4)=a_4-2b_{4,3}+c_{3,2}-2d_5, & \phi_*(a_5)=a_5+2d_3, & \phi_*(a_6)=a_6+2d_2,\\ \phi_*(a_7)=a_7, & \phi_*(a_8)=a_8. & \end{array}\right. \tag{5.3}$$

5.3. Calculations for the pair $(\mathfrak{D}_1,\mathfrak{D}_2)$. — In this part we provide the last two steps of the calculations.

Step 1 – We have isomorphisms

$$W(\mathfrak{D}_1)\simeq W(D_1)\oplus W\big(A\cdot D_{\widehat{P}_2}\big)\oplus W\big(AfA\cdot D_{\widehat{P}_2}\big),\qquad W(\mathfrak{D}_2)\simeq W\big(D_{\widehat{P}_1}\big)\oplus W(AD_2)\oplus W\big(AfA\cdot D_{\widehat{P}_2}\big).$$

We transport the bases of $W(D_2)$ and $W(D_{\widehat{P}_2})$ by A and $A\phi A$. Therefore, if we take algebraic bases of $W\big(D_{\widehat{P}_1}\big)$, $W\big(D_{\widehat{P}_2}\big)$ and $W(D_2)$, the matrix of $\psi_*\,:\,W(\mathfrak{D}_1)\longrightarrow W(\mathfrak{D}_2)$ has the form

	$A\phi\cdot(\mathfrak{l}_i)_{1\leq i\leq 10}$	$A\phi\cdot(A\mathfrak{m}_i)_{1\leq i\leq 10}$	$A\phi\cdot(A\phi A\mathfrak{m}_i)_{1\leq i\leq 10}$
$(\mathfrak{l}_i)_{1\leq i\leq 10}$	$0_{10\times 10}$	$0_{10\times 10}$	$\mathscr{Q}$
$(A\mathfrak{m}_i)_{1\leq i\leq 10}$ $\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2(5A\Delta)\big)^{\dagger}$	$\mathscr{L}\mathscr{K}^{-1}$	$0_{55\times 10}$	$0_{55\times 10}$
$(A\phi A\mathfrak{m}_i)_{1\leq i\leq 10}$	$0_{10\times 10}$	$\mathrm{Id}_{10\times 10}$	$0_{10\times 10}$

– *Matrix* $\mathscr{Y}$ –

The matrix $\mathscr{Q}$ is the 10×10 matrix given explicitly by

$$\begin{bmatrix}
-4\lambda_1-1 & q_{1,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
-4 & 4\mu_1-1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
8-25\lambda_2 & \mu_1(25\lambda_2-8) & 6\lambda_2-1 & q_{3,4} & 0 & 0 & 0 & 0 & 0 & 0\\
-25 & 25\mu_1 & 6 & 1-6\mu_2 & 0 & 0 & 0 & 0 & 0 & 0\\
-50\lambda_3 & 8+50\lambda_3\mu_1 & 4(1+3\lambda_3) & -4\mu_2(1+3\lambda_3) & -1 & \mu_3+\lambda_3 & 0 & 0 & 0 & 0\\
-50 & 50\mu_1 & 12 & -12\mu_2 & 0 & 1 & 0 & 0 & 0 & 0\\
q_{7,1} & q_{7,2} & 18\lambda_4-42 & \mu_2(42-18\lambda_4) & -10 & 10\mu_3-10 & -1 & \mu_4+\lambda_4 & 0 & 0\\
q_{8,1} & q_{8,2} & 18 & -18\mu_2 & 0 & 0 & 0 & 1 & 0 & 0\\
q_{9,1} & q_{9,2} & q_{9,3} & q_{9,4} & -110 & 110\mu_3-62 & -12 & 12\mu_4 & -1 & q_{9,10}\\
q_{10,1} & q_{10,2} & 24 & -24\mu_2 & 0 & 0 & 0 & 0 & 0 & 1
\end{bmatrix}$$

where

$$\begin{aligned}
&q_{1,2}=\mu_1(1+4\lambda_1)-\lambda_1, \quad q_{3,4}=\mu_2(1-6\lambda_2)+\lambda_2, \quad q_{7,1}=2\alpha+200+\lambda_4(3\mu_4^2-\lambda_4^2-75-4\lambda_4)+2\mu_4^3,\\
&q_{7,2}=40+\mu_1\big(75\lambda_4-2\alpha-200+4\lambda_4^2-2\mu_4^3-3\mu_4^2\lambda_4+\lambda_4^3\big)-(\mu_4+\lambda_4)^2, \quad q_{8,1}=3(\mu_4^2-\lambda_4^2)-75-8\lambda_4,\\
&q_{8,2}=\mu_1\big(75+8\lambda_4+3(\lambda_4^2-\mu_4^2)\big)-2(\mu_4+\lambda_4), \quad q_{9,1}=672-4\mu_5(\lambda_5+1)+24(\mu_4^3+\alpha)-2(\lambda_5^2+\mu_5^2)-100\lambda_5,\\
&q_{9,2}=\mu_1\big(4\mu_5(1+\lambda_5)-24(\mu_4^3+\alpha)+2(\mu_5^2+\lambda_5^2)+100\lambda_5-672\big)+320-2\alpha-12\mu_4^2, \quad q_{9,3}=24\lambda_5-92-4\alpha,\\
&q_{9,4}=\mu_2(92+4\alpha-24\lambda_5), \quad q_{9,10}=\mu_5+1+\lambda_5, \quad q_{10,1}=-4(25+\lambda_5+\mu_5), \quad q_{10,2}=4\mu_1(25+\lambda_5+\mu_5).
\end{aligned}$$

Step 2 – We have $V(\mathfrak{D}_1)=\mathfrak{h}(\mathbb{P}^2)$ and $V(\mathfrak{D}_2)=\mathrm{H}^0\big(\mathbb{P}^2,\mathrm{T}\mathbb{P}^2\big(5A\Delta\big)\big)$. Let us decompose a basis of $V(\mathfrak{D}_1)$ in an algebraic basis of $W(\mathfrak{D}_1)$:

	$(a_i)_{1\leq i\leq 8}$
$(\mathfrak{l}_i)_{1\leq i\leq 10}$	$\mathscr{M}_{\mathrm{a}}$
$(A\mathfrak{m}_i)_{1\leq i\leq 10}$	$\mathscr{M}'_{\mathrm{a}}$
$(A\phi A\mathfrak{m}_i)_{1\leq i\leq 10}$	$\mathscr{M}''_{\mathrm{a}}$

– *Matrix* $\mathscr{V}_1$ –

To compute quickly the matrices $\mathscr{M}'_{\mathrm{a}}$ and $\mathscr{M}''_{\mathrm{a}}$, we make the following remark (which can be proved by an easy computation): for any holomorphic vector field defined in a neighborhood $(0,0)$, if we lift it as a section of $W(D_{\widehat{P}_1})$ or $W\big(D_{\widehat{P}_2}\big)$, this section depends only on the seven Taylor components ∂_y, $y\partial_y$, $z\partial_y$,∂_z, $y\partial_z$, $z\partial_z$, $y^2\partial_z$ of the vector field at $(0,0)$. The corresponding sections in algebraic bases are given by the tables

	∂_y	$y\partial_y$	$z\partial_y$	∂_z	$y\partial_z$	$z\partial_z$	$y^2\partial_z$
$\mathfrak{l}_1$	$-\lambda_1$	0	0	1	0	0	0
$\mathfrak{l}_2$	-1	0	0	0	0	0	0
$\mathfrak{l}_3$	2	0	0	0	$-\lambda_2$	0	0
$\mathfrak{l}_4$	0	0	0	0	-1	0	0
$\mathfrak{l}_5$	0	0	0	0	$-2\lambda_3$	0	0
$\mathfrak{l}_6$	0	0	0	0	-2	0	0
$\mathfrak{l}_7$	$-\lambda_4^2$	-3	0	λ_4^3	$-3\lambda_4$	2	0
$\mathfrak{l}_8$	$-2\lambda_4$	0	0	$3\lambda_4^2$	-3	0	0
$\mathfrak{l}_9$	0	0	-3	$2\lambda_5^2$	$-4\lambda_5$	0	-2
$\mathfrak{l}_{10}$	0	0	0	$4\lambda_5$	-4	0	0

– *Matrix* $\mathscr{Z}_1$ –

	∂_y	$y\partial_y$	$z\partial_y$	∂_z	$y\partial_z$	$z\partial_z$	$y^2\partial_z$
$\mathfrak{m}_1$	$-\mu_1$	0	0	1	0	0	0
$\mathfrak{m}_2$	-1	0	0	0	0	0	0
$\mathfrak{m}_3$	2	0	0	0	$-\mu_2$	0	0
$\mathfrak{m}_4$	0	0	0	0	-1	0	0
$\mathfrak{m}_5$	0	0	0	0	$-2\mu_3$	0	0
$\mathfrak{m}_6$	0	0	0	0	-2	0	0
$\mathfrak{m}_7$	$-\mu_4^2$	3	0	$-\mu_4^3$	$-3\mu_4$	-2	0
$\mathfrak{m}_8$	$-2\mu_4$	0	0	$-3\mu_4^2$	-3	0	0
$\mathfrak{m}_9$	0	0	3	$2\mu_5^2$	$-4\mu_5$	0	-2
$\mathfrak{m}_{10}$	0	0	0	$4\mu_5$	-4	0	0

– *Matrix* $\mathscr{Z}_2$ –

Therefore, in order to compute $\mathscr{M}'_{\mathrm{a}}$ and $\mathscr{M}''_{\mathrm{a}}$, it suffices to extract the seven aforementioned Taylor components of the vector field $A_*^{-1}a_i$ $\big($resp. $(A\phi A)_*^{-1}a_i\big)$ at $(0,0)$. Then we multiply the resulting vector by the matrix $\mathscr{Z}_1$ $\big($resp. $\mathscr{Z}_2\big)$ and we obtain $\mathscr{M}'_{\mathrm{a}}$ $\big($resp. $\mathscr{M}''_{\mathrm{a}}\big)$. We won't give the exact expressions of $\mathscr{M}'_{\mathrm{a}}$ and $\mathscr{M}''_{\mathrm{a}}$ because of lack of space.

We now deal with $V(\mathfrak{D}_2)$. We decompose a basis of vectors of $V(\mathfrak{D}_2)$ in an algebraic basis of $W(\mathfrak{D}_2)$. We get the following matrix

	$(Aa_i)_{1\leq i\leq 8}$	$(Ab_{p,q})_{0\leq p\leq q+1\leq m+1}$	$(Ac_{p,q})_{0\leq p\leq q+1\leq m+1}$	$(Ad_p)_{1\leq p\leq m}$
$(\mathfrak{l}_i)_{1\leq i\leq 10}$	$\mathscr{N}'_{\mathrm{a}}$	$\mathscr{N}'_{\mathrm{b}}$	$\mathscr{N}'_{\mathrm{c}}$	$\mathscr{N}'_{\mathrm{d}}$
$(A\mathfrak{m}_i)_{1\leq i\leq 10}$	$\mathscr{N}_{\mathrm{a}}$	$\mathscr{N}_{\mathrm{b}}$	$\mathscr{N}_{\mathrm{c}}$	$\mathscr{N}_{\mathrm{d}}$
$(Ab_{p,q})_{0\leq p\leq q+1\leq m+1}$	$0_{25\times 8}$	$\mathrm{Id}_{25\times 25}$	$0_{25\times 25}$	$0_{25\times 25}$
$(Ac_{p,q})_{0\leq p\leq q+1\leq m+1}$	$0_{25\times 8}$	$0_{25\times 25}$	$\mathrm{Id}_{25\times 25}$	$0_{25\times 25}$
$(Ad_p)_{1\leq p\leq m}$	$0_{5\times 8}$	$0_{5\times 25}$	$0_{5\times 25}$	$\mathrm{Id}_{5\times 5}$
$(A\phi A\mathfrak{m}_i)_{1\leq i\leq 10}$	$\mathscr{N}''_{\mathrm{a}}$	$\mathscr{N}''_{\mathrm{b}}$	$\mathscr{N}''_{\mathrm{c}}$	$\mathscr{N}''_{\mathrm{d}}$

– *Matrix* $\mathscr{V}_2$ –

The matrices $\mathscr{N}'_{\mathrm{a}}$ $\big($resp. $\mathscr{N}''_{\mathrm{a}}\big)$, $\mathscr{N}'_{\mathrm{b}}$ $\big($resp. $\mathscr{N}''_{\mathrm{b}}\big)$, $\mathscr{N}'_{\mathrm{c}}$ $\big($resp. $\mathscr{N}''_{\mathrm{c}}\big)$ and $\mathscr{N}'_{\mathrm{d}}$ $\big($resp. $\mathscr{N}''_{\mathrm{d}}\big)$ appearing in $\mathscr{V}_2$ are computed in the same way as $\mathscr{N}'_{\mathrm{a}}$ $\big($resp. $\mathscr{N}''_{\mathrm{a}}\big)$.

5.4. The result. — We can now state and prove the following result.

Theorem 5.1. — *Let X be the rational surface obtained by blowing up the projective plane 15 times at the infinitely near points $\widehat{P}_1$, $A\widehat{P}_2$ and $A\phi A\widehat{P}_2$ and ψ be the lift of $A\phi$ as an automorphism of X. Then the characteristic polynomial Q_{ψ} of the map ψ_* acting on the space $\mathrm{H}^1\big(X, \mathrm{T}X\big)$ of infinitesimal deformations of X is*

$$Q_{\psi}(x) = \big(x^2+3x+1\big)\,\big(x^2+18x+1\big)\,\big(x^2-7x+1\big)\,\big(x^2+x+1\big)\,\big(x-1\big)^2\,\big(x+1\big)^4\,\big(x^2-x+1\big)^4.$$

Besides, there is only one nontrivial Jordan block, which is a 2×2 Jordan block attached with the eigenvalue -1.

Proof. — Let $\mathscr{E}_1$ be the subspace of $W(\mathfrak{D}_1)$ of dimension 22 defined by

$$\mathscr{E}_1 = \mathrm{Span}\left\{\begin{array}{c} \mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3, \mathfrak{l}_4, \mathfrak{l}_5, \mathfrak{l}_6, \mathfrak{l}_7, \mathfrak{l}_8, \mathfrak{l}_9, \mathfrak{l}_{10}, \\ A\mathfrak{m}_1, A\mathfrak{m}_2, A\mathfrak{m}_3, A\mathfrak{m}_4, A\mathfrak{m}_5, A\mathfrak{m}_6, A\mathfrak{m}_7, \\ A\phi A\mathfrak{m}_1, A\phi A\mathfrak{m}_2, A\phi A\mathfrak{m}_3, A\phi A\mathfrak{m}_4, A\phi A\mathfrak{m}_5 \end{array}\right\}.$$

A direct calculation shows that $\mathscr{E}_1$ is a direct factor of $V(\mathfrak{D}_1)$ in $W(\mathfrak{D}_1)$. If $\mathscr{E}_2$ is the image of $\mathscr{E}_1$ in $W(\mathfrak{D}_2)$ by the natural injection, then $\mathscr{E}_2$ is a direct factor of $V(\mathfrak{D}_2)$ in $W(\mathfrak{D}_2)$.

Therefore, the composition of the morphisms (5.2) can be expressed as

$$\mathscr{E}_1 \xrightarrow{\;j\;} \mathscr{E}_1 \oplus V(\mathfrak{D}_1) \simeq W(\mathfrak{D}_1) \xrightarrow{\;\psi_*\;} W(\mathfrak{D}_2) \simeq \mathscr{E}_2 \oplus V(\mathfrak{D}_2) \xrightarrow{\;p\;} \mathscr{E}_2 \simeq \mathscr{E}_1 \ .$$

The matrix of j (resp. p) can be computed using $\mathscr{V}_1$ $\big($resp. $\mathscr{V}_2\big)$ and the matrix of ψ_* is $\mathscr{Y}$. □

Corollary 5.2. — *If $(\mathfrak{X}, \mathfrak{f})$ is a holomorphic family of rational surfaces endowed with an automorphism that lifts the pair (X, ψ), then $\mathfrak{m}(\mathfrak{X}) \leq 2$.*

It is an interesting problem to know if this bound for $\mathfrak{m}(\mathfrak{X})$ is sharp.

References

[1] M. F. Atiyah and R. Bott. A Lefschetz fixed point formula for elliptic complexes. II. Applications. *Ann. of Math. (2)*, 88:451–491, 1968.

[2] E. Bedford and K. Kim. Continuous families of rational surface automorphisms with positive entropy. *Math. Ann.*, 348(3):667–688, 2010.

[3] J. Blanc. Dynamical degrees of (pseudo)-automorphisms fixing cubic hypersurfaces, `arxiv:1204.4256`. *Indiana Univ. Math. J.*, to appear.

[4] J. Blanc and J. Déserti. Embeddings of $\mathrm{SL}(2,\mathbb{Z})$ into the Cremona group. *Transform. Groups*, 17(1):21–50, 2012.

[5] M. Brunella. *Birational geometry of foliations*. Monografías de Matemática. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2000. Available electronically at http://www.impa.br/Publicacoes/Monografias/Abstracts/brunella.ps.

[6] S. Cantat and C. Favre. Symétries birationnelles des surfaces feuilletées. *J. Reine Angew. Math.*, 561:199–235, 2003.

[7] J. Déserti and J. Grivaux. Special automorphisms of rational surfaces with positive topological entropy, `arxiv:1004.0656`. *Indiana Univ. Math. J.*, to appear.

[8] J. Diller. Cremona transformations, surface automorphisms, and plane cubics. *Michigan Math. J.*, 60(2):409–440, 2011. With an appendix by I. Dolgachev.

[9] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1994. Reprint of the 1978 original.

[10] A. Lins Neto. Some examples for the Poincaré and Painlevé problems. *Ann. Sci. École Norm. Sup. (4)*, 35(2):231–266, 2002.

[11] C. T. McMullen. Dynamics on blowups of the projective plane. *Publ. Math. Inst. Hautes Études Sci.*, (105):49–89, 2007.

[12] H. Poincaré. Sur l'intégration algébrique des équations différentielles du premier ordre et du premier degré. *Rend. Circ. Mat. Palermo*, 5:161–191, 1891.

[13] H. Poincaré. Sur l'intégration algébriques des équations différentielles. *C. R. Math. Acad. Sci. Paris*, 112:761–764, 1891.

[14] H. Poincaré. Sur l'intégration algébrique des équations différentielles du premier ordre et du premier degré. *Rend. Circ. Mat. Palermo*, 11:193–239, 1897.

[15] L. Puchuri Medina. Degree of the first integral of a foliation in the pencil $\mathcal{P}_4$, `arxiv:1105.0104`.

6 Maple computations

Contents

6 Maple computations **1**
6.1 **First definitions** 1
6.2 **The procedures to compute direct images of vector fields** 2
6.2.1 The procedures `champ` 2
6.2.2 The procedures `devlim` and `extract` 3
6.3 **Definition of the residues matrices** 10
6.4 **Construction of the matrix $\mathscr{Y}$** 13
6.4.1 Construction of a block matrix 13
6.4.2 Computation of $\mathscr{Q}$ 15
6.5 **The 8 vectors of $V(\mathfrak{D}_1)$ that form the columns of the matrix $\mathscr{V}_1$** 17
6.5.1 Columns of $\mathscr{M}_a$ (eight (10×1)-vectors) 17
6.5.2 Columns of $\mathscr{M}'_a$ (eight (10×1)-vectors) 17
6.5.3 Columns of $\mathscr{M}''_a$ (eight (10×1)-vectors) 18
6.5.4 Construction of the matrix $\mathscr{V}_1$ 19
6.6 **The 63 vectors of $V(\mathfrak{D}_2)$ that form the columns of the matrix $\mathscr{V}_2$** 19
6.6.1 Vectors of type a (we compute the columns of the matrices $\mathscr{N}'_{\mathrm{a}}$, $\mathscr{N}_{\mathrm{a}}$ et $\mathscr{N}''_{\mathrm{a}}$) 19
6.6.2 Vectors of type b (we compute the columns of the matrices $\mathscr{N}'_{\mathrm{b}}$, $\mathscr{N}_{\mathrm{b}}$ and $\mathscr{N}''_{\mathrm{b}}$) 22
6.6.3 Vectors of type c (we compute the columns of the matrices $\mathscr{N}'_{\mathrm{c}}$, $\mathscr{N}_{\mathrm{c}}$ and $\mathscr{N}''_{\mathrm{c}}$) 28
6.6.4 Vectors of type d (we compute the columns of the matrices $\mathscr{N}'_{\mathrm{d}}$, $\mathscr{N}_{\mathrm{d}}$ and $\mathscr{N}''_{\mathrm{d}}$) 34
6.6.5 Construction of the matrix $\mathscr{V}_2$ 36
6.7 **Computation of the action of ψ_* on $\mathrm{H}^1(X, \mathrm{T}X)$** 36

```
>  restart:
>  with(LinearAlgebra):
```

6.1 First definitions

```
>  restart:
>  with(LinearAlgebra):
```

The function `f` is the function ϕ of the text given in the affine coordinates (y, z).

```
>  f1:=(y,z)->(y*z^2)/(z^2+y^3):
>  f2:=(y,z)->(z^3)/(z^2+y^3):
>  f:=(y,z)->(f1(y,z),f2(y,z)):
```

The inverse of `f` is denoted by `g`.

```
>  g1:=(y,z)->y*z^2/(z^2-y^3):
>  g2:=(y,z)->z^3/(z^2-y^3):
>  g:=(y,z)->(g1(y,z),g2(y,z)):
```

We define the action of the matrix `A` in the coordinates (y, z), its inverse is `B`.

```
>  A:=(y,z)->((-1+(1+alpha)*z)/(alpha+2*(1-alpha)*y+(2+alpha-alpha^2)*z),(1-2*y+(1-alpha)*
z)/(alpha+2*(1-alpha)*y+(2+alpha-alpha^2)*z)):
>  B:=(y,z)->((1-y-(1+alpha)*z)/(1+alpha+(alpha-3)*y+(1-alpha^2)*z),(1+y+(1-alpha)*z)/(1+alpha+
(alpha-3)*y+(1-alpha^2)*z)):
```

We define $\mathtt{m} = \mathtt{AfAfA}$ and $\mathtt{n} = \mathtt{BgBgB}$.

```
>  m1:=(y,z)->op(1,[A(f(A(f(A(y,z)))))]):
>  m2:=(y,z)->op(2,[A(f(A(f(A(y,z)))))]):
>  m:=(y,z)->(m1(y,z), m2(y,z)):
>  n1:=(y,z)->op(1,[B(g(B(g(B(y,z)))))]):
>  n2:=(y,z)->op(2,[B(g(B(g(B(y,z)))))]):
>  n:=(y,z)->(n1(y,z), n2(y,z)):
```

6.2 The procedures to compute direct images of vector fields

6.2.1 The procedures `champ`

The procedure `champ1` associates to a vector field Z the coordinates of the Taylor expansion of the vector field $A_* Z$ in the following order: ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z, $y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$. The variable of the procedure is not the vector field itself but a curve $t \mapsto M_t$ whose the derivate at 0 is Z. For example, if $Z = u(y,z)\partial_y + v(y,z)\partial_z$, we can take $M_t(y,z) = (u(y,z) + ty, v(y,z) + tz)$.

```
>  champ1:=proc(M)
>  local prod,res,res1,res2,u,v:
>  global resu1,resu2,w:
>  prod:=unapply([(A@(M@B))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=[subs({y=0,z=0},res1(y,z)),subs({y=0,z=0},diff(res1(y,z),y)),subs({y=0,z=0},diff(
res1(y,z),z))]:
>  resu2:=[subs({y=0,z=0},res2(y,z)),subs({y=0,z=0},diff(res2(y,z),y)),subs({y=0,z=0},diff(
res2(y,z),z)),subs({y=0,z=0},diff(diff(res2(y,z),y),y)/2)]:
>  u:=op(resu1):v:=op(resu2):
>  w:=[u,v]:
>  RETURN(convert(w,Vector))
>  end:
```

The procedure `champ2` associates to a vector field Z the coordinates of the Taylor expansion of the vector field $(\phi A)_*^{-1}\,Z$ in the following order: ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z, $y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$.

```
>  champ2:=proc(M)
>  local prod,res,res1,res2,u,v:
>  global resu1,resu2,w:
>  prod:=unapply([(B@(g@(M@(f@A))))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=[subs({y=0,z=0},res1(y,z)),subs({y=0,z=0},diff(res1(y,z),y)),subs({y=0,z=0},diff(
res1(y,z),z))]:
>  resu2:=[subs({y=0,z=0},res2(y,z)),subs({y=0,z=0},diff(res2(y,z),y)),subs({y=0,z=0},diff(
res2(y,z),z)),subs({y=0,z=0},diff(diff(res2(y,z),y),y)/2)]:
>  u:=op(resu1):v:=op(resu2):
>  w:=[u,v]:
>  RETURN(convert(w,Vector))
>  end:
```

The procedure `champ3` associates to a vector field Z the coordinates of the Taylor expansion of the vector field $(A\phi A)_*^{-1}\,(Z)$ in the following order: ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z, $y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$.

```
>  champ3:=proc(M)
>  local prod,res,res1,res2:
>  global resu1,resu2:
>  prod:=unapply([(B@(g@(B@(M@(A@(f@A))))))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=[subs({y=0,z=0},res1(y,z)),subs({y=0,z=0},diff(res1(y,z),y)),subs({y=0,z=0},diff(
res1(y,z),z))]:
>  resu2:=[subs({y=0,z=0},res2(y,z)),subs({y=0,z=0},diff(res2(y,z),y)),subs({y=0,z=0},diff(
res2(y,z),z)),subs({y=0,z=0},diff(diff(res2(y,z),y),y)/2)]:
>  RETURN(Transpose(convert([seq(factor(op(k,resu1)),k=1..nops(resu1)),seq(factor(op(k,resu2)
),k=1..nops(resu2))],Matrix))):
>  end:
```

The procedure `champ4` associates to a vector field Z the coordinates of the Taylor expansion of the vector field $A_*^{-1}\,(Z)$ in the following order: ∂_y, $y\partial_y$, $z\partial_y$, ∂_z, $y\partial_z$, $z\partial_z$, $y^2\partial_z$.

```
>  champ4:=proc(M)
>  local prod,res,res1,res2,u,v:
>  global resu1,resu2,w:
>  prod:=unapply([(B@(M@A))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
```

```
>  resu1:=[subs({y=0,z=0},res1(y,z)),subs({y=0,z=0},diff(res1(y,z),y)),subs({y=0,z=0},diff(
res1(y,z),z))]:
>  resu2:=[subs({y=0,z=0},res2(y,z)),subs({y=0,z=0},diff(res2(y,z),y)),subs({y=0,z=0},diff(
res2(y,z),z)),subs({y=0,z=0},diff(diff(res2(y,z),y),y)/2)]:
>  u:=op(resu1):v:=op(resu2):
>  w:=[u,v]:
>  RETURN(convert(w,Vector))
>  end:
```

The procedure `champ5` associates to a vector field Z the coordinates of the Taylor expansion of the vector field $(A\phi A\phi A)_*^{-1}(Z)$ in the following order: ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z, $y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$.

```
>  champ5:=proc(M)
>  local prod,res,res1,res2,u,v:
>  global resu1,resu2,w:
>  prod:=unapply([(m@(M@n))(y,z)],t):
>  res:=(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=[subs({y=0,z=0},res1(y,z)),subs({y=0,z=0},diff(res1(y,z),y)),subs({y=0,z=0},diff(
res1(y,z),z))]:
>  resu2:=[subs({y=0,z=0},res2(y,z)),subs({y=0,z=0},diff(res2(y,z),y)),subs({y=0,z=0},diff(
res2(y,z),z)),subs({y=0,z=0},diff(diff(res2(y,z),y),y)/2)]:
>  u:=op(resu1):v:=op(resu2):
>  w:=[u,v]:
>  RETURN(convert(w,Vector))
>  end:
```

This procedure is too heavy for Maple, we will not use it. To replace it, we will do successive Taylor expansions at each step of blow-ups. We deal with it in the next section.

6.2.2 The procedures `devlim` and `extract`

We compute the points where we take the Taylor expansions.

```
>  A(0,0);
```

$-\alpha^{-1}, \alpha^{-1}$

```
>  simplify(op(1,[(A@f@A)(0,0)])), simplify(op(2,[(A@f@A)(0,0)]));
```

α^{-1}, α^{-1}

```
>  simplify(op(1,[(A@f@A@f@A)(0,0)])), simplify(op(2,[(A@f@A@f@A)(0,0)]));
```

$0, 0$

The procedure `devlim1` associates to a vector field Z in coordinates (y, z) (still given by the derivative at 0 of a curve M_t) the Taylor expansion of A_*Z of order 2 at a point (p, q). The result is viewed as a polynomial vector field in a neighborhood of (p, q) and the procedure returns a curve N_t associated to this vector field.

```
>  devlim1:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(y,z)->(op(M(y,z))):
>  prod:=unapply([(A@(MMM@B))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=simplify(mtaylor(res1(y,z),[y=p,z=q],3)):
>  resu2:=simplify(mtaylor(res2(y,z),[y=p,z=q],3)):
>  N:=unapply([y+t*resu1,z+t*resu2],(y,z)):
>  RETURN(N):
>  end:
```

The procedure `devlim2` is similar to devlim1 but computes the Taylor expansion of $(A\phi)_*Z$ instead of the Taylor expansion of A_*Z.

```
>  devlim2:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(y,z)->(op(M(y,z))):
>  prod:=unapply([(A@f@(MMM@g@B))(y,z)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  resu1:=simplify(mtaylor(res1(y,z),[y=p,z=q],3)):
```

```
>  resu2:=simplify(mtaylor(res2(y,z),[y=p,z=q],3)):
>  N:=unapply([y+t*resu1,z+t*resu2],(y,z)):
>  RETURN(N):
>  end:
```

The procedure `extract` associates to a vector field given by a curve $N_t(y,z)$ the components in ∂_y, $y\partial_y$, $z\partial_y$, ∂_z, $y\partial_z$, $z\partial_z$ and $y^2\partial_z$ in the Taylor expansion of the associated vector field at $(0,0)$.

```
>  extract:=proc(M)
>  local prod,res,res1,res2,w:
>  prod:=unapply(M(y,z),t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(y,z)):
>  res2:=unapply(op(2,res),(y,z)):
>  w[1]:=simplify(res1(0,0)):
>  w[2]:=simplify(subs({y=0,z=0},diff(res1(y,z),y))):
>  w[3]:=simplify(subs({y=0,z=0},diff(res1(y,z),z))):
>  w[4]:=simplify(res2(0,0)):
>  w[5]:=simplify(subs({y=0,z=0},diff(res2(y,z),y))):
>  w[6]:=simplify(subs({y=0,z=0},diff(res2(y,z),z))):
>  w[7]:=simplify(subs({y=0,z=0},diff(res2(y,z),y$2)/2)):
>  w:=[w[1],w[2],w[3],w[4],w[5],w[6],w[7]]:
>  RETURN(convert(w,Vector)):
>  end:
```

The function $\texttt{psi}_1$ is the projection of the blow-up of $(0,0)_{y,z}$, and $\texttt{eta}_1$ is its inverse.

```
>  psi1:=(u,v)->(u,u*v):
>  eta1:=(y,z)->(y,z/y):
```

The function $\texttt{psi}_2$ is the projection of the blow-up of $(-1/\alpha,1/\alpha)_{y,z}$ and $\texttt{eta}_2$ is its inverse.

```
>  psi2:=(u,v)->(u-1/alpha,u*v+1/alpha):
>  eta2:=(y,z)->(y+1/alpha,(z-1/alpha)/(y+1/alpha)):
```

The function $\texttt{psi}_3$ is the projection of the blow-up of $(1/\alpha,1/\alpha)_{y,z}$ and $\texttt{eta}_3$ is its inverse.

```
>  psi3:=(u,v)->(u+1/alpha, u*v+1/alpha):
>  eta3:=(y,z)->(y-1/alpha,(z-1/alpha)/(y-1/alpha)):
```

The procedure `u1devlim1` associates to a vector field Z on $\mathbb{P}^2$ blown up at $(0,0)_{y,z}$ written in the coordinates (u_1,v_1) the Taylor expansion of A_*Z of order 3 at a point (p,q). The result is considered as a polynomial vector field in a neighborhood of (p,q) in the coordinates (u_1,v_1) which are this time the coordinates on $\mathbb{P}^2$ blown up at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, and the procedure gives a curve N_t associated to this vector field.

```
>  u1devlim1:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(u,v)->(op(M(u,v))):
>  prod:=unapply([(eta2@(A@(psi1@MMM@eta1@(B@psi2))))(u,v)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(u,v)):
>  res2:=unapply(simplify(op(2,res)),(u,v)):
>  resu1:=simplify(mtaylor(res1(u,v),[u=p,v=q],4)):
>  resu2:=simplify(mtaylor(res2(u,v),[u=p,v=q],4)):
>  N:=unapply([u+t*resu1,v+t*resu2],(u,v)):
>  RETURN(N):
>  end:
```

We compute the points where we take the Taylor expansions.

```
>  limit(op(1,[(eta2@(A@psi1))(u,v)]),{u=0,v=0}), limit(op(2,[(eta2@(A@psi1))(u,v)]),{u=0,
v=0});
```

$0,\ -(1-\alpha)^{-1}$

```
>  limit(op(1,[(eta3@(A@f@A)@psi1)(u,v)]),{u=0,v=0}), limit(op(2,[(eta3@(A@f@A)@psi1)(u,v)
]),{u=0,v=0});
```

$0,\ -(-1-\alpha)^{-1}$

```
>  limit(op(1,[(eta1@(A@f@A@f@A)@psi1)(u,v)]),{u=0,v=0}), limit(op(2,[(eta1@(A@f@A@f@A)@psi1)
(u,v)]),{u=0,v=0});
```

$0,\ 0$

The procedure **u1devlim2** associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ expressed in the coordinates (u_1,v_1) the Taylor expansion of $(A\phi)_*Z$ of order 3 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (u_1,v_1) which are this time some coordinates on $\mathbb{P}^2$ blown up at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, and the procedure returns a curve N_t associated to this vector field.

```
>  u1devlim2:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(u,v)->(op(M(u,v))):
>  prod:=unapply([(eta3@(A@f@(psi2@MMM@eta2@g@B@psi3)))(u,v)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(u,v)):
>  res2:=unapply(simplify(op(2,res)),(u,v)):
>  resu1:=simplify(mtaylor(res1(u,v),[u=p,v=q],4)):
>  resu2:=simplify(mtaylor(res2(u,v),[u=p,v=q],4)):
>  N:=unapply([u+t*resu1,v+t*resu2],(u,v)):
>  RETURN(N):
>  end:
```

The procedure **u1devlim3** associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ expressed in the coordinates (u_1,v_1) the Taylor expansion of $(A\phi)_*Z$ of order 3 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (u_1,v_1) which are this time coordinates on $\mathbb{P}^2$ blown up at $(0,0)_{y,z}$, and the procedure gives a curve N_t associated to this vector field.

```
>  u1devlim3:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(u,v)->(op(M(u,v))):
>  prod:=unapply([(eta1@(A@f@(psi3@MMM@eta3@g@B@psi1)))(u,v)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(u,v)):
>  res2:=unapply(simplify(op(2,res)),(u,v)):
>  resu1:=simplify(mtaylor(res1(u,v),[u=p,v=q],4)):
>  resu2:=simplify(mtaylor(res2(u,v),[u=p,v=q],4)):
>  N:=unapply([u+t*resu1,v+t*resu2],(u,v)):
>  RETURN(N):
>  end:
```

The procedure **u1extract** associates to a vector field given by a curve $N_t(u_1,v_1)$ the components in ∂_{u_1}, $u_1\,\partial_{u_1}$, $v_1\,\partial_{u_1}$, $u_1v_1\,\partial_{u_1}$, $v_1^2\,\partial_{u_1}$, $v_1^3\,\partial_{u_1}$, ∂_{v_1}, $u_1\,\partial_{v_1}$, $v_1\,\partial_{v_1}$ and $v_1^2\,\partial_{v_1}$ of the Taylor expansion of the associated vector field at $(0,0)$.

```
>  u1extract:=proc(M)
>  local prod,res,res1,res2,w:
>  prod:=unapply(M(u,v),t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(u,v)):
>  res2:=unapply(op(2,res),(u,v)):
>  w[1]:=simplify(res1(0,0)):
>  w[2]:=simplify(subs({u=0,v=0},diff(res1(u,v),u))):
>  w[3]:=simplify(subs({u=0,v=0},diff(res1(u,v),v))):
>  w[4]:=simplify(subs({u=0,v=0},diff(res1(u,v),u,v))):
>  w[5]:=simplify(subs({u=0,v=0},diff(res1(u,v),v$2))/2):
>  w[6]:=simplify(subs({u=0,v=0},diff(res1(u,v),v$3))/3):
>  w[7]:=simplify(res2(0,0)):
>  w[8]:=simplify(subs({u=0,v=0},diff(res2(u,v),u))):
>  w[9]:=simplify(subs({u=0,v=0},diff(res2(u,v),v))):
>  w[10]:=simplify(subs({u=0,v=0},diff(res2(u,v),v$2)/2)):
>  w:=[w[1],w[2],w[3],w[4],w[5],w[6],w[7],w[8],w[9],w[10]]:
>  RETURN(convert(w,Vector)):
>  end:
```

The function $\texttt{ppsi}_1$ is the projection of the blow up of $(0,0)_{y,z}$ and $(0,0)_{u_1,v_1}$, and $\texttt{eeta}_1$ is its inverse.

```
>  ppsi1:=(r,s)->(r*s,r*s^2):
>  eeta1:=(y,z)->(y^2/z,z/y):
```

The function $\texttt{ppsi}_2$ is the projection of the blow up of $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$, and $\texttt{eeta}_2$ is its inverse.

```
>  ppsi2:=(r,s)->(r*s-1/alpha, r*s*(s+1/(alpha-1))+1/alpha):
>  eeta2:=(y,z)->((y+1/alpha)/((z-1/alpha)/(y+1/alpha)-1/(alpha-1)),(z-1/alpha)/(y+1/alpha)
-1/(alpha-1)):
```

The function `ppsi`$_3$ is the projection of the blow up of $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$, and `eeta`$_3$ is its inverse.

```
>  ppsi3:=(r,s)->(r*s+1/alpha, r*s*(s+1/(alpha+1))+1/alpha):
>  eeta3:=(y,z)->((y-1/alpha)/((z-1/alpha)/(y-1/alpha)-1/(alpha+1)),(z-1/alpha)/(y-1/alpha)
-1/(alpha+1)):
```

We compute the points where we take the Taylor expansions.

```
>  limit(op(1,[(eeta2@(A@ppsi1))(r,s)]),{r=0,s=0}), limit(op(2,[(eeta2@(A@ppsi1))(r,s)]),{r=0,
s=0});

0, 0
>  limit(op(1,[(eeta3@(A@f@A)@ppsi1)(r,s)]),{r=0,s=0}), limit(op(2,[(eeta3@(A@f@A)@ppsi1)(
r,s)]),{r=0,s=0});

0, 0
>  limit(op(1,[(eeta1@(A@f@A@f@A)@ppsi1)(r,s)]),{r=0,s=0}), limit(op(2,[(eeta1@(A@f@A@f@A)
@ppsi1)(r,s)]),{r=0,s=0});

0, 0
```

The procedure `r2devlim1` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $(0,0)_{y,z}$ and $(0,0)_{u_1,v_1}$ expressed in coordinates (r_2,s_2) the Taylor expansion of A_*Z of order 2 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_2,s_2) which are this time coordinates on $\mathbb{P}^2$ blown up at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$, and the procedure returns a curve N_t associated to this vector field.

```
>  r2devlim1:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeta2@(A@(ppsi1@MMM@eeta1@(B@ppsi2))))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],3)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],3)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure `r2devlim2` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$ expressed in coordinates (r_2,s_2) the Taylor expansion of $(A\phi)_*\,Z$ of order 2 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_2,s_2) which are this time the coordinates of $\mathbb{P}^2$ blown up at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$, and the procedure returns a curve N_t associated to this vector field.

```
>  r2devlim2:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeta3@(A@f@(ppsi2@MMM@eeta2@g@B@ppsi3)))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],3)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],3)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure `r2devlim3` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$ and $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$ expressed in coordinates (r_2,s_2) the Taylor expansion of $(A\phi)_*Z$ of order 2 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_2,s_2) which are this time the coordinates of $\mathbb{P}^2$ blown up at $(0,0)_{y,z}$ and $(0,0)_{u_1,v_1}$, and the procedure returns a curve N_t associated to this vector field.

```
>  r2devlim3:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeta1@(A@f@(ppsi3@MMM@eeta3@g@B@ppsi1)))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
```

```
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],3)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],3)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure `r2extract` associates to a vector field given by a curve $N_t(r_2, s_2)$ the components in ∂_{r_2}, $r_2\,\partial_{r_2}$, $s_2\,\partial_{r_2}$, $r_2 s_2\,\partial_{r_2}$, $r_2^2\,\partial_{r_2}$, $s_2^2\,\partial_{r_2}$, ∂_{s_2}, $r_2\,\partial_{s_2}$, $s_2\,\partial_{s_2}$, $r_2 s_2\,\partial_{s_2}$, $r_2^2\,\partial_{s_2}$ and $s_2^2\,\partial_{s_2}$ in the Taylor expansion of the associated vector field at $(0,0)$.

```
>  r2extract:=proc(M)
>  local prod,res,res1,res2,w:
>  prod:=unapply(M(r,s),t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(r,s)):
>  res2:=unapply(op(2,res),(r,s)):
>  w[1]:=simplify(res1(0,0)):
>  w[2]:=simplify(subs({r=0,s=0},diff(res1(r,s),r))):
>  w[3]:=simplify(subs({r=0,s=0},diff(res1(r,s),s))):
>  w[4]:=simplify(subs({r=0,s=0},diff(res1(r,s),r,s))):
>  w[5]:=simplify(subs({r=0,s=0},diff(res1(r,s),r$2))/2):
>  w[6]:=simplify(subs({r=0,s=0},diff(res1(r,s),s$2))/2):
>  w[7]:=simplify(res2(0,0)):
>  w[8]:=simplify(subs({r=0,s=0},diff(res2(r,s),r))):
>  w[9]:=simplify(subs({r=0,s=0},diff(res2(r,s),s))):
>  w[10]:=simplify(subs({r=0,s=0},diff(res2(r,s),r,s))):
>  w[11]:=simplify(subs({r=0,s=0},diff(res2(r,s),r$2))/2):
>  w[12]:=simplify(subs({r=0,s=0},diff(res2(r,s),s$2))/2):
>  w:=[w[1],w[2],w[3],w[4],w[5],w[6],w[7],w[8],w[9],w[10],w[11],w[12]]:
>  RETURN(convert(w,Vector)):
>  end:
```

The function $\texttt{pppsi}_1$ is the projection of the blow up of $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and $\texttt{eeeta}_1$ is its inverse.

```
>  pppsi1:=(r,s)->(r*s*s,r*s*s^2):
>  eeeta1:=(y,z)->((y^2/z)/(z/y),(z/y)):
```

The function $\texttt{pppsi}_2$ is the projection of the blow up of $\left(-\frac{1}{\alpha}, \frac{1}{\alpha}\right)_{y,z}$, $\left(0, \frac{1}{\alpha-1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and $\texttt{eeeta}_2$ is its inverse.

```
>  pppsi2:=(r,s)->(r*s*s-1/alpha, r*s*s*(s+1/(alpha-1))+1/alpha):
>  eeeta2:=(y,z)->(((y+1/alpha)/((z-1/alpha)/(y+1/alpha)-1/(alpha-1)))/((z-1/alpha)/(y+1/alpha)
-1/(alpha-1)),((z-1/alpha)/(y+1/alpha)-1/(alpha-1))):
```

The function $\texttt{pppsi}_3$ is the projection of the blow up of $\left(\frac{1}{\alpha}, \frac{1}{\alpha}\right)_{y,z}$, $\left(0, \frac{1}{\alpha+1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and $\texttt{eeeta}_3$ is its inverse.

```
>  pppsi3:=(r,s)->(r*s*s+1/alpha, r*s*s*(s+1/(alpha+1))+1/alpha):
>  eeeta3:=(y,z)->(((y-1/alpha)/((z-1/alpha)/(y-1/alpha)-1/(alpha+1)))/((z-1/alpha)/(y-1/alpha)
-1/(alpha+1)),((z-1/alpha)/(y-1/alpha)-1/(alpha+1))):
```

We compute the points where we take the Taylor expansions.

```
>  limit(op(1,[(eeeta2@(A@pppsi1))(r,s)]),{r=1,s=0}), limit(op(2,[(eeeta2@(A@pppsi1))(r,s)
]),{r=1,s=0});
```

$$\frac{-(\alpha-1)^5}{2\alpha^4},\ 0$$

```
>  limit(op(1,[(eeeta3@(A@f@A)@pppsi1)(r,s)]),{r=1,s=0}), limit(op(2,[(eeeta3@(A@f@A)@pppsi1)
(r,s)]),{r=1,s=0});
```

$$\frac{(1+\alpha)^5}{2\alpha^4},\ 0$$

```
>  limit(op(1,[(eeeta1@(A@f@A@f@A)@pppsi1)(r,s)]),{r=1,s=0}), limit(op(2,[(eeeta1@(A@f@A@f@A)
@pppsi1)(r,s)]),{r=1,s=0});
```

$$-1,\ 0$$

The procedure `r3devlim1` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$ expressed in coordinates (r_3,s_3) the Taylor expansion of A_*Z of order 1 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_3,s_3) which are this time the coordinates on $\mathbb{P}^2$ blown up at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and the procedure gives a curve N_t associated to this vector field.

```
>   r3devlim1:=proc(M,p,q)
>   local prod,res,res1,res2,resu1,resu2,N,MMM:
>   MMM:=(r,s)->(op(M(r,s))):
>   prod:=unapply([(eeeta2@(A@(pppsi1@MMM@eeeta1@(B@pppsi2))))(r,s)],t):
>   res:=simplify(subs(t=0,diff(prod(t),t))):
>   res1:=unapply(simplify(op(1,res)),(r,s)):
>   res2:=unapply(simplify(op(2,res)),(r,s)):
>   resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],2)):
>   resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],2)):
>   N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>   RETURN(N):
>   end:
```

The procedure `r3devlim2` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$ expressed in the coordinates (r_3,s_3) the Taylor expansion of $(A\phi)_*Z$ of order 1 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_3,s_3) which are this time the coordinates of $\mathbb{P}^2$ blown up at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and the procedure returns a curve N_t associated to this vector field.

```
>   r3devlim2:=proc(M,p,q)
>   local prod,res,res1,res2,resu1,resu2,N,MMM:
>   MMM:=(r,s)->(op(M(r,s))):
>   prod:=unapply([(eeeta3@(A@f@(pppsi2@MMM@eeeta2@g@B@pppsi3)))(r,s)],t):
>   res:=simplify(subs(t=0,diff(prod(t),t))):
>   res1:=unapply(simplify(op(1,res)),(r,s)):
>   res2:=unapply(simplify(op(2,res)),(r,s)):
>   resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],2)):
>   resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],2)):
>   N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>   RETURN(N):
>   end:
```

The procedure `r3devlim3` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$ expressed in coordinates (r_3,s_3) the Taylor expansion of $(A\phi)_*Z$ of order 1 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_3,s_3) which are this time the coordinates on $\mathbb{P}^2$ blown up at $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$ and $(0,0)_{r_2,s_2}$, and the procedure returns a curve N_t associated to this vector field.

```
>   r3devlim3:=proc(M,p,q)
>   local prod,res,res1,res2,resu1,resu2,N,MMM:
>   MMM:=(r,s)->(op(M(r,s))):
>   prod:=unapply([(eeeta1@(A@f@(pppsi3@MMM@eeeta3@g@B@pppsi1)))(r,s)],t):
>   res:=simplify(subs(t=0,diff(prod(t),t))):
>   res1:=unapply(simplify(op(1,res)),(r,s)):
>   res2:=unapply(simplify(op(2,res)),(r,s)):
>   resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],2)):
>   resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],2)):
>   N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>   RETURN(N):
>   end:
```

The procedure `r3extract` associates to a vector field given by a curve $N_t(r_3,s_3)$ the components of ∂_{r_3}, $s_3\partial_{r_3}$ and ∂_{s_3} in the Taylor expansion of the associated vector field at $(0,0)$.

```
>   r3extract:=proc(M)
>   local prod,res,res1,res2,w:
>   prod:=unapply(M(r,s),t):
>   res:=simplify(subs(t=0,diff(prod(t),t))):
>   res1:=unapply(op(1,res),(r,s)):
>   res2:=unapply(op(2,res),(r,s)):
>   w[1]:=simplify(res1(-1,0)):
>   w[2]:=simplify(subs({r=-1,s=0},diff(res1(r,s),s))):
>   w[3]:=simplify(res2(-1,0)):
>   w:=[w[1],w[2],w[3]]:
>   RETURN(convert(w,Vector)):
>   end:
```

The function `ppppsi`$_1$ is the projection of the blow up of $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$ $(0,0)_{r_2,s_2}$ and $(1,0)_{r_3,s_3}$, and `eeeeta`$_1$ is its inverse.

```
>  ppppsi1:=(r,s)->((r*s+1)*s*s,(r*s+1)*s*s^2):
>  eeeeta1:=(y,z)->(((y^2/z)/(z/y)-1)/(z/y),(z/y)):
```

The function `ppppsi`$_2$ is the projection of the blow up at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(-\frac{(\alpha-1)^5}{2\alpha^4},0\right)_{r_3,s_3}$, and `eeeeta`$_2$ is its inverse.

```
>  ppppsi2:=(r,s)->((r*s-(1/2)*(-1+alpha)^5/alpha^4)*s*s-1/alpha, (r*s-(1/2)*(-1+alpha)^5/
alpha^4)*s*s*(s+1/(alpha-1))+1/alpha):
>  eeeeta2:=(y,z)->((((y+1/alpha)/((z-1/alpha)/(y+1/alpha)-1/(alpha-1)))/((z-1/alpha)/(y+1/
alpha)-1/(alpha-1))+(1/2)*(-1+alpha)^5/alpha^4)/((z-1/alpha)/(y+1/alpha)-1/(alpha-1)),((z-1/
alpha)/(y+1/alpha)-1/(alpha-1))):
```

The function `ppppsi`$_3$ is the projection of the blow up at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(\frac{(\alpha+1)^5}{2\alpha^4},0\right)$, and `eeeeta`$_3$ is its inverse.

```
>  ppppsi3:=(r,s)->((r*s+(1/2)*(1+alpha)^5/alpha^4)*s*s+1/alpha, (r*s+(1/2)*(1+alpha)^5/alpha^4)
*s*s*(s+1/(alpha+1))+1/alpha):
>  eeeeta3:=(y,z)->((((y-1/alpha)/((z-1/alpha)/(y-1/alpha)-1/(alpha+1)))/((z-1/alpha)/(y-1/
alpha)-1/(alpha+1))-(1/2)*(1+alpha)^5/alpha^4)/((z-1/alpha)/(y-1/alpha)-1/(alpha+1)),((z-1/
alpha)/(y-1/alpha)-1/(alpha+1))):
```

The function `ppppsi`$_4$ is the projection of the blow up at $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$ $(0,0)_{r_2,s_2}$ and $(-1,0)_{r_3,s_3}$, and `eeeeta`$_1$ is its inverse.

```
>  ppppsi4:=(r,s)->((r*s-1)*s*s,(r*s-1)*s*s^2):
>  eeeeta4:=(y,z)->(((y^2/z)/(z/y)+1)/(z/y),(z/y)):
```

We compute the points where we take the Taylor expansions.

```
>  limit(op(1,[(eeeeta2@(A@ppppsi1))(r,s)]),{r=0,s=0}), limit(op(2,[(eeeeta2@(A@ppppsi1))(
r,s)]),{r=0,s=0});
```

$$\frac{3(\alpha-1)^4(1-\alpha^2+\alpha^3-\alpha)}{4\alpha^5},\ 0$$

```
>  limit(op(1,[(eeeeta3@(A@f@A)@ppppsi1)(r,s)]),{r=0,s=0}), limit(op(2,[(eeeeta3@(A@f@A)@ppppsi1)
(r,s)]),{r=0,s=0});
```

$$\frac{-3(1+\alpha)^4(-\alpha-1+\alpha^2+\alpha^3)}{4\alpha^5},\ 0$$

```
>  limit(op(1,[(eeeeta4@(A@f@A@f@A)@ppppsi1)(r,s)]),{r=0,s=0}), limit(op(2,[(eeeeta4@(A@f@A@f@A)
@ppppsi1)(r,s)]),{r=0,s=0});
```

$$0,\ 0$$

The procedure `r4devlim1` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $(1,0)_{r_3,s_3}$ expressed in the coordinates (r_4,s_4) the Taylor expansion of A_*Z of order 0 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_4,s_4) which are this time the coordinates on $\mathbb{P}^2$ blown up at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(-\frac{(\alpha-1)^5}{2\alpha^4},0\right)_{r_3,s_3}$, and the procedure returns a curve N_t associated to this vector field.

```
>  r4devlim1:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeeeta2@(A@(ppppsi1@MMM@eeeeta1@(B@ppppsi2))))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],1)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],1)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure `r4devlim2` associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(-\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha-1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(-\frac{(\alpha-1)^5}{2\alpha^4},0\right)_{r_3,s_3}$ expressed in the coordinates (r_4,s_4) the Taylor expansion of $(A\phi)_*\,Z$ of order

0 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_4,s_4) which are this time the coordinates on $\mathbb{P}^2$ blown up at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(\frac{(\alpha+1)^5}{2\alpha^4},0\right)$, and the procedure returns a curve N_t associated to this vector field.

```
>  r4devlim2:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeeeta3@(A@f@(ppppsi2@MMM@eeeeta2@g@B@ppppsi3)))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],1)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],1)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure **r4devlim3** associates to a vector field Z on the blow up of $\mathbb{P}^2$ at $\left(\frac{1}{\alpha},\frac{1}{\alpha}\right)_{y,z}$, $\left(0,\frac{1}{\alpha+1}\right)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $\left(\frac{(\alpha+1)^5}{2\alpha^4},0\right)$, and expressed in coordinates (r_4,s_4) the Taylor expansion of $(A\phi)_*\,Z$ of order 0 at a point (p,q). The result is viewed as a polynomial vector field in a neighborhood of (p,q) in the coordinates (r_4,s_4) which are this time the coordinates on $\mathbb{P}^2$ blown up at $(0,0)_{y,z}$, $(0,0)_{u_1,v_1}$, $(0,0)_{r_2,s_2}$ and $(-1,0)_{r_3,s_3}$, and the procedure returns a curve N_t associated to this vector field.

```
>  r4devlim3:=proc(M,p,q)
>  local prod,res,res1,res2,resu1,resu2,N,MMM:
>  MMM:=(r,s)->(op(M(r,s))):
>  prod:=unapply([(eeeeta4@(A@f@(ppppsi3@MMM@eeeeta3@g@B@ppppsi4)))(r,s)],t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(simplify(op(1,res)),(r,s)):
>  res2:=unapply(simplify(op(2,res)),(r,s)):
>  resu1:=simplify(mtaylor(res1(r,s),[r=p,s=q],1)):
>  resu2:=simplify(mtaylor(res2(r,s),[r=p,s=q],1)):
>  N:=unapply([r+t*resu1,s+t*resu2],(r,s)):
>  RETURN(N):
>  end:
```

The procedure **r4extract** associates to a vector field given by a curve $N_t(r_4,s_4)$ the components of ∂_{r_4} and ∂_{s_4} in the Taylor expansion of the vector field associated at $(0,0)$.

```
>  r4extract:=proc(M)
>  local prod,res,res1,res2,w:
>  prod:=unapply(M(r,s),t):
>  res:=simplify(subs(t=0,diff(prod(t),t))):
>  res1:=unapply(op(1,res),(r,s)):
>  res2:=unapply(op(2,res),(r,s)):
>  w[1]:=simplify(res1(0,0)):
>  w[2]:=simplify(res2(0,0)):
>  w:=[w[1],w[2]]:
>  RETURN(convert(w,Vector)):
>  end:
```

6.3 Definition of the residues matrices

$\mathtt{R}_1$, denoted by $\mathscr{X}_1$ in the article, is the (10×7)-matrix of ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z,$y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  R_1:=Transpose(Matrix([[-lambda[1],-1,2,0,0,0,-lambda[4]^2,-2*lambda[4],0,0],[0,0,0,0,0,
0,-3,0,0,0],[0,0,0,0,0,0,0,0,-3,0],[1,0,0,0,0,0,lambda[4]^3,3*lambda[4]^2,2*lambda[5]^2,4*
lambda[5]],[0,0,-lambda[2],-1,-2*lambda[3],-2,-3*lambda[4],-3,-4*lambda[5],-4],[0,0,0,0,0,
0,2,0,0,0],[0,0,0,0,0,0,0,0,-2,0]]));
```

$$R\_1 := \begin{bmatrix} -\lambda_1 & 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & -\lambda_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2\,\lambda_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 \\ -{\lambda_4}^2 & -3 & 0 & {\lambda_4}^3 & -3\,\lambda_4 & 2 & 0 \\ -2\,\lambda_4 & 0 & 0 & 3\,{\lambda_4}^2 & -3 & 0 & 0 \\ 0 & 0 & -3 & 2\,{\lambda_5}^2 & -4\,\lambda_5 & 0 & -2 \\ 0 & 0 & 0 & 4\,\lambda_5 & -4 & 0 & 0 \end{bmatrix}$$

$\mathtt{R}_2$, denoted by $\mathscr{L}_2$ in the article, is the (10×7)-matrix of ∂_y, $y\,\partial_y$, $z\,\partial_y$, ∂_z,$y\,\partial_z$, $z\,\partial_z$, $y^2\,\partial_z$ in the basis $(\mathfrak{m}_i)_{1\le i\le 10}$.

```
> R_2:=Transpose(Matrix([[-mu[1],-1,2,0,0,0,-mu[4]^2,-2*mu[4],0,0],[0,0,0,0,0,0,3,0,0,0],
[0,0,0,0,0,0,0,0,3,0],[1,0,0,0,0,0,-mu[4]^3,-3*mu[4]^2,2*mu[5]^2,4*mu[5]],[0,0,-mu[2],-1,-2*
mu[3],-2,-3*mu[4],-3,-4*mu[5],-4],[0,0,0,0,0,0,-2,0,0,0],[0,0,0,0,0,0,0,0,-2,0]]));
```

$$R\_2 := \begin{bmatrix} -\mu_1 & 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & -\mu_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2\,\mu_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 \\ -{\mu_4}^2 & 3 & 0 & -{\mu_4}^3 & -3\,\mu_4 & -2 & 0 \\ -2\,\mu_4 & 0 & 0 & -3\,{\mu_4}^2 & -3 & 0 & 0 \\ 0 & 0 & 3 & 2\,{\mu_5}^2 & -4\,\mu_5 & 0 & -2 \\ 0 & 0 & 0 & 4\,\mu_5 & -4 & 0 & 0 \end{bmatrix}$$

$\mathtt{RR}_1$ is the (10×10)-matrix that gives the residues of ∂_{u_1}, $u_1\,\partial_{u_1}$, $v_1\,\partial_{u_1}$, $u_1v_1\,\partial_{u_1}$, $v_1^2\,\partial_{u_1}$, $v_1^3\,\partial_{u_1}$, ∂_{v_1}, $u_1\,\partial_{v_1}$, $v_1\,\partial_{v_1}$, $v_1^2\,\partial_{v_1}$ in the basis $(\mathfrak{l}_i)_{1\le i\le 10}$.

```
> RR_1:=Matrix([[0,0,0,0,0,0,0,0,0,0],[0,0,0,0,0,0,0,0,0,0],[1,0,0,0,0,0,-lambda[2],0,0,0],
[0,0,0,0,0,0,-1,0,0,0],[0,0,1,0,0,0,-2*lambda[3],0,0,0],[0,0,0,0,0,0,-2,0,0,0],[0,-1,0,0,1,
0,-3*lambda[4],0,2,0],[0,0,0,0,0,0,-3,0,0,0],[0,0,0,-1,0,1,-4*lambda[5],-2,0,2],[0,0,0,0,0,
0,-4,0,0,0]]);
```

$$RR\_1 := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & -\lambda_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & -2\,\lambda_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 & 0 & -3\,\lambda_4 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 & -4\,\lambda_5 & -2 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & -4 & 0 & 0 & 0 \end{bmatrix}$$

$\mathtt{RRR}_1$ is the (10×12)-matrix that gives the residues of ∂_{r_2}, $r_2\,\partial_{r_2}$, $s_2\,\partial_{r_2}$, $r_2s_2\,\partial_{r_2}$, $r_2^2\partial_{r_2}$, $s_2^2\,\partial_{r_2}$, ∂_{s_2}, $r_2\,\partial_{s_2}$, $s_2\,\partial_{s_2}$, $r_2s_2\,\partial_{s_2}$, $r_2^2\,\partial_{s_2}$, $s_2^2\,\partial_{s_2}$ in the basis $(\mathfrak{l}_i)_{1\le i\le 10}$.

```
> RRR_1:=Matrix([[0,0,0,0,0,0,0,0,0,0,0,0],[0,0,0,0,0,0,0,0,0,0,0,0],[0,0,0,0,0,0,0,0,0,0,
0,0],[0,0,0,0,0,0,0,0,0,0,0,0],[1,0,0,0,0,0,-lambda[3],0,0,0,0,0],[0,0,0,0,0,0,-1,0,0,0,0,
0],[0,-1,1,0,0,0,-2*lambda[4],-1,1,0,0,0],[0,0,0,0,0,0,-2,0,0,0,0,0],[0,0,0,-1,1,1,-3*lambda[5],
0,0,-1,1,1],[0,0,0,0,0,0,-3,0,0,0,0,0]]);
```

$$RRR\_1 := \begin{bmatrix} \text{` 10 x 12 `}\,(\mathit{Matrix}) \\ \text{`Data Type: `}\ \mathit{anything} \\ \text{`Storage: `}\ \mathit{rectangular} \\ \text{`Order: `}\ \mathit{Fortran\_order} \end{bmatrix}$$

```
> evalm(RRR_1);
```

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & -\lambda_3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & -2\,\lambda_4 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 1 & -3\,\lambda_5 & 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

`RRRR`$_1$ is the (10×3)-matrix that gives the residues of ∂_{r_3}, $s_3\partial_{r_3}$, ∂_{s_3} in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
> RRRR_1:=Matrix([[0,0,0],[0,0,0],[0,0,0],[0,0,0],[0,0,0],[0,0,0],[1,0,-lambda[4]],[0,0,-1],
[0,1,-2*lambda[5]],[0,0,-2]]);
```

$$RRRR\_1 := \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & -\lambda_4 \\ 0 & 0 & -1 \\ 0 & 1 & -2\,\lambda_5 \\ 0 & 0 & -2 \end{bmatrix}$$

`RRRRR`$_1$ is the (10×2)-matrix that gives the residues of ∂_{r_4}, ∂_{s_4} in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
> RRRRR_1:=Matrix([[0,0],[0,0],[0,0],[0,0],[0,0],[0,0],[0,0],[0,0],[1,-lambda[5]],[0,-1]])
;
```

$$RRRRR\_1 := \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 1 & -\lambda_5 \\ 0 & -1 \end{bmatrix}$$

`P` is the transition matrix from the algebraic basis $(\mathfrak{m}_i)_{1\leq 10}$ to the geometric basis for $\widehat{P}_2$.

```
>  P:=Matrix([[-mu[1],1,0,0,0,0,0,0,0,0],[-1,0,0,0,0,0,0,0,0,0],[2,0,1,-mu[2],0,0,0,0,0,0],
[0,0,0,-1,0,0,0,0,0,0],[0,0,0,-2*mu[3],1,-mu[3],0,0,0,0],[0,0,0,-2,0,-1,0,0,0,0],[-mu[4]^2,
-mu[4]^3,0,-3*mu[4],0,-2*mu[4],1,-mu[4],0,0],[-2*mu[4],-3*mu[4]^2,0,-3,0,-2,0,-1,0,0],[0,2*
mu[5]^2,0,-4*mu[5],0,-3*mu[5],0,-2*mu[5],1,-mu[5]],[0,4*mu[5],0,-4,0,-3,0,-2,0,-1]]);
```

$$P := \begin{bmatrix} -\mu_1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 1 & -\mu_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2\,\mu_3 & 1 & -\mu_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -{\mu_4}^2 & -{\mu_4}^3 & 0 & -3\,\mu_4 & 0 & -2\,\mu_4 & 1 & -\mu_4 & 0 & 0 \\ -2\,\mu_4 & -3\,{\mu_4}^2 & 0 & -3 & 0 & -2 & 0 & -1 & 0 & 0 \\ 0 & 2\,{\mu_5}^2 & 0 & -4\,\mu_5 & 0 & -3\,\mu_5 & 0 & -2\,\mu_5 & 1 & -\mu_5 \\ 0 & 4\,\mu_5 & 0 & -4 & 0 & -3 & 0 & -2 & 0 & -1 \end{bmatrix}$$

`PP`, denoted by $\mathscr{K}$ in the article, is the transition matrix from the algebraic basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$ to the geometric basis $(\mathfrak{e}_i)_{1\leq i\leq 10}$ for $\widehat{P}_1$.

```
>  PP:=Matrix([[-lambda[1],1,0,0,0,0,0,0,0,0],[-1,0,0,0,0,0,0,0,0,0],[2,0,1,-lambda[2],0,0,
0,0,0,0],[0,0,0,-1,0,0,0,0,0,0],[0,0,0,-2*lambda[3],1,-lambda[3],0,0,0,0],[0,0,0,-2,0,-1,0,
0,0,0],[-lambda[4]^2, lambda[4]^3,0,-3*lambda[4],0,-2*lambda[4],1,-lambda[4],0,0],[-2*lambda[4],
3*lambda[4]^2,0,-3,0,-2,0,-1,0,0],[0,2*lambda[5]^2,0,-4*lambda[5],0,-3*lambda[5],0,-2*lambda[5],
1,-lambda[5]],[0,4*lambda[5],0,-4,0,-3,0,-2,0,-1]]);
```

$$PP := \begin{bmatrix} -\lambda_1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 1 & -\lambda_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2\,\lambda_3 & 1 & -\lambda_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -{\lambda_4}^2 & {\lambda_4}^3 & 0 & -3\,\lambda_4 & 0 & -2\,\lambda_4 & 1 & -\lambda_4 & 0 & 0 \\ -2\,\lambda_4 & 3\,{\lambda_4}^2 & 0 & -3 & 0 & -2 & 0 & -1 & 0 & 0 \\ 0 & 2\,{\lambda_5}^2 & 0 & -4\,\lambda_5 & 0 & -3\,\lambda_5 & 0 & -2\,\lambda_5 & 1 & -\lambda_5 \\ 0 & 4\,\lambda_5 & 0 & -4 & 0 & -3 & 0 & -2 & 0 & -1 \end{bmatrix}$$

`PPP` is the transition matrix from the algebraic basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$ to the geometric basis $(\mathfrak{e}_i)_{1\leq i\leq 10}$ for $\widehat{P}_1$.

```
>  PPP:=MatrixInverse(PP):
```

6.4 Construction of the matrix $\mathscr{Y}$

6.4.1 Construction of a block matrix

We input the coefficients of the matrix $\mathscr{L}$.

```
>  col[1]:=[-mu[1], -1, 2, 0, 0, 0, 2*mu[4]^2, 4*mu[4], 0, 0, 0, 0, 0, 0, 0, 0, -4, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 3, 0];
```

$$col_1 := [-\mu_1, -1, 2, 0, 0, 0, 2\,\mu_4^2, 4\,\mu_4, 0, 0, 0, 0, 0, 0, 0, 0, -4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 3, 0]$$

```
>  col[2]:=
>  [1, 0, 0, 0, 0, 0, -mu[4]^3, -3*mu[4]^2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 2, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, -2];
```

$$col_2 := [1,0,0,0,0,0,-\mu_4^3,-3\,\mu_4^2,0,0,0,0,0,0,0,0,0,0,0,0,0,2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-2]$$

```
>  col[3]:=
>  [0, 0, 1, 0, 0, 0, mu[4]^2, 2*mu[4], 0, 0, 0, 0, 0, 0, 0, 0, -2, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 1, 0];
```

$$col_3 := [0,0,1,0,0,0,\mu_4^2,2\,\mu_4,0,0,0,0,0,0,0,0,-2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0]$$

```
>  col[4]:=
>  [0, 0, -mu[2], -1, 2*mu[3]^2-2*mu[3], 4*mu[3]-2, mu[4], 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 2, 0, 0];
```

$$col_4 := [0,0,-\mu_2,-1,2\,\mu_3^2-2\,\mu_3,4\,\mu_3-2,\mu_4,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,2,0,0]$$

```
>  col[5]:=
>  [0, 0, -2*mu[2], -2, (mu[3]-1)^2, 2*(mu[3]-1), 0, 0, 0, 0, 0, 0, -2, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0];
```

$$col_5 := [0,0,-2\,\mu_2,-2,(\mu_3-1)^2,2\,\mu_3-2,0,0,0,0,0,0,-2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0]$$

```
>  col[6]:=
>  [0, 0, 0, 0, mu[3]^2-mu[3], 2*mu[3]-1, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0];
```

$$col_6 := [0,0,0,0,\mu_3^2-\mu_3,2\,\mu_3-1,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0]$$

```
>  col[7]:=
>  [0, 0, mu[2]^2, 2*mu[2], 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 1, 0, 0, 0];
```

$$col_7 := [0,0,\mu_2^2,2\,\mu_2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0]$$

```
>  col[8]:=
>  [0, 0, mu[2], 1, 0, 0, 0, 0, 0, 0, 0, 0, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0];
```

$$col_8 := [0,0,\mu_2,1,0,0,0,0,0,0,0,0,2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

```
>  col[9]:=[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 1, 0, 0, 0, 0];
```

$$col_9 := [0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0]$$

```
>  col[10]:=
>  [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
0, 0, 0, 0, 0];
```

$$col_{10} := [0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

The matrix `PART` is by definition $\mathscr{L}\mathscr{K}^{-1}$.

```
>  PART:=MatrixMatrixMultiply(Transpose(Matrix([col[1],col[2],col[3],col[4],col[5],col[6],
col[7],col[8],col[9],col[10]])),PPP);
```

$$PART := \left[\begin{array}{c} \texttt{‘ 65 x 10 ‘}\ (\mathit{Matrix}) \\ \texttt{‘Data Type: }\ \text{‘ }\mathit{anything} \\ \texttt{‘Storage: }\ \text{‘ }\mathit{rectangular} \\ \texttt{‘Order: }\ \text{‘ }\mathit{Fortran\_order} \end{array}\right]$$

```
>  for k from 1 to 85 do
>  for j from 1 to 30 do
>  L[k][j]:=0
>  od
>  od:
```

We construct the matrix $\mathscr{Y}$, whose upper right block $\mathscr{Q}$ is indeterminate.

```
>  for k from 1 to 10 do
>  for l from 1 to 10 do
>  L[k][l+20]:=Q[k,l]
>  od
>  od:
>  for k from 76 to 85 do
>  L[k][k-65]:=1
>  od:
>  for k from 11 to 75 do
>  for j from 1 to 10 do
>  L[k][j]:=PART[k-10,j]
>  od
>  od:
>  for k from 1 to 85 do
>  L[k]:=convert(L[k],list)
>  od:
>  MM:=Matrix([seq(L[k],k=1..85)]):
```

6.4.2 Computation of $\mathscr{Q}$

To compute the coefficients of $\mathscr{Q}$, we consider the matrix `R` obtained by multiplying $\mathscr{Q}$ at right by the transition matrix from the algebraic basis to the geometric basis of $\widehat{P}_2$. The matrix `R` is then the matrix

$$(A\phi A\phi A)_* : W(\widehat{P}_2, \mathscr{B}_{\texttt{geom}}) \longrightarrow W(\widehat{P}_1, \mathscr{B}_{\texttt{alg}}).$$

Computation of $(A\phi A\phi A)_*\, \partial_y$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  y:='y':  z:='z':
>  M:=(y,z)->[y+t,z]:
>  U:=MatrixMatrixMultiply(R_1,extract(devlim2(devlim2((devlim1(M,-1/alpha,1/alpha)), 1/alpha,
1/alpha),0,0))):
>  for k from 1 to 10 do
>  R[k][1]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_*\, \partial_z$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  y:='y':  z:='z':
>  M:=(y,z)->[y,z+t]:
>  U:=MatrixMatrixMultiply(R_1,extract(devlim2(devlim2((devlim1(M,-1/alpha,1/alpha)), 1/alpha,
1/alpha),0,0))):
>  for k from 1 to 10 do
>  R[k][2]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_*\, \partial_{u_1}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  u:='u':  v:='v':
>  M:=(u,v)->[u+t,v]:
>  U:=MatrixMatrixMultiply(RR_1,u1extract(u1devlim3(u1devlim2(u1devlim1(M,0,1/(alpha-1)), 0,
1/(1+alpha)),0,0))):
>  for k from 1 to 10 do
>  R[k][3]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_*\, \partial_{v_1}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  u:='u':  v:='v':
>  M:=(u,v)->[u,v+t]:
>  U:=MatrixMatrixMultiply(RR_1,u1extract(u1devlim3(u1devlim2(u1devlim1(M,0,1/(alpha-1)), 0,
1/(1+alpha)),0,0))):
>  for k from 1 to 10 do
>  R[k][4]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_*\, \partial_{r_2}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r+t,s]:
>  U:=MatrixMatrixMultiply(RRR_1,r2extract(r2devlim3(r2devlim2(r2devlim1(M,0,0),0,0),0,0))
):
```

```
>  for k from 1 to 10 do
>  R[k][5]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_* \, \partial_{s_2}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r,s+t]:
>  U:=MatrixMatrixMultiply(RRR_1,r2extract(r2devlim3(r2devlim2(r2devlim1(M,0,0),0,0),0,0))
):
>  for k from 1 to 10 do
>  R[k][6]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_* \, \partial_{r_3}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r+t,s]:
>  U:=MatrixMatrixMultiply(RRRR_1,r3extract(r3devlim3(r3devlim2(r3devlim1(M,-(1/2)*(-1+alpha)
^5/alpha^4,0),(1/2)*(1+alpha)^5/alpha^4, 0),-1,0))):
>  for k from 1 to 10 do
>  R[k][7]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_* \, \partial_{s_3}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r,s+t]:
>  U:=MatrixMatrixMultiply(RRRR_1,r3extract(r3devlim3(r3devlim2(r3devlim1(M,-(1/2)*(-1+alpha)
^5/alpha^4,0),(1/2)*(1+alpha)^5/alpha^4, 0),-1,0))):
>  for k from 1 to 10 do
>  R[k][8]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_* \, \partial_{c_4}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r+t,s]:
>  U:=MatrixMatrixMultiply(RRRRR_1,r4extract(r4devlim3(r4devlim2(r4devlim1(M,(3/4)*(-1+alpha)
^4*(alpha^3-alpha-alpha^2+1)/alpha^5,0),-(3/4)*(1+alpha)^4*(-alpha-1+alpha^3+alpha^2)/alpha^5,
0),0,0))):
>  for k from 1 to 10 do
>  R[k][9]:=U[k];
>  od:
```

Computation of $(A\phi A\phi A)_* \, \partial_{d_4}$ in the basis $(\mathfrak{l}_i)_{1\leq i\leq 10}$.

```
>  r:='r':  s:='s':
>  M:=(r,s)->[r,s+t]:
>  U:=MatrixMatrixMultiply(RRRRR_1,r4extract(r4devlim3(r4devlim2(r4devlim1(M,(3/4)*(-1+alpha)
^4*(alpha^3-alpha-alpha^2+1)/alpha^5,0),-(3/4)*(1+alpha)^4*(-alpha-1+alpha^3+alpha^2)/alpha^5,
0),0,0))):
>  for k from 1 to 10 do
>  R[k][10]:=U[k];
>  od:
```

We construct the matrix `R`, then we compute $\mathscr{Q}$.

```
>  for k from 1 to 10 do
>  R[k]:=convert(R[k],list)
>  od:
>  TEMP:=Matrix([seq(R[k],k=1..10)]):
>  Q:=MatrixMatrixMultiply(TEMP, MatrixInverse(P)):
```

We can display the coefficients of $\mathscr{Q}$.

```
>  simplify(Q[9,4]);
```

$$4\,(23+\alpha-6\,\lambda_5)\mu_2$$

6.5 The 8 vectors of $V(\mathfrak{D}_1)$ that form the columns of the matrix $\mathscr{V}_1$

6.5.1 Columns of $\mathscr{M}_a$ (eight (10×1)-vectors)

a_1

```
> pro1:=[-lambda[1],-1,2,0,0,0,-lambda[4]^2,-2*lambda[4],0,0];
> a[1]:=convert(pro1,Vector):
```

$\mathit{pro1} := [-\lambda_1, -1, 2, 0, 0, 0, -{\lambda_4}^2, -2\,\lambda_4, 0, 0]$

a_2

```
> pro2:=[0,0,0,0,0,0,-3,0,0,0];
> a[2]:=convert(pro2,Vector):
```

$\mathit{pro2} := [0, 0, 0, 0, 0, 0, -3, 0, 0, 0]$

a_3

```
> pro3:=[0,0,0,0,0,0,0,0,-3,0];
> a[3]:=convert(pro3,Vector):
```

$\mathit{pro3} := [0, 0, 0, 0, 0, 0, 0, 0, -3, 0]$

a_4

```
> pro4:=[1,0,0,0,0,0,lambda[4]^3,3*lambda[4]^2,2*lambda[5]^2,4*lambda[5]];
> a[4]:=convert(pro4,Vector):
```

$\mathit{pro4} := [1, 0, 0, 0, 0, 0, {\lambda_4}^3, 3\,{\lambda_4}^2, 2\,{\lambda_5}^2, 4\,\lambda_5]$

a_5

```
> pro5:=[0,0,-lambda[2],-1,-2*lambda[3],-2,-3*lambda[4],-3,-4*lambda[5],-4];
> a[5]:=convert(pro5,Vector):
```

$\mathit{pro5} := [0, 0, -\lambda_2, -1, -2\,\lambda_3, -2, -3\,\lambda_4, -3, -4\,\lambda_5, -4]$

a_6

```
> pro6:=[0,0,0,0,0,0,2,0,0,0];
> a[6]:=convert(pro6,Vector):
```

$\mathit{pro6} := [0, 0, 0, 0, 0, 0, 2, 0, 0, 0]$

a_7

```
> pro7:=[0,0,0,0,0,0,0,0,0,0];
> a[7]:=convert(pro7,Vector):
```

$\mathit{pro7} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

a_8

```
> pro8:=[0,0,0,0,0,0,0,0,0,0];
> a[8]:=convert(pro8,Vector):
```

$\mathit{pro8} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

6.5.2 Columns of $\mathscr{M}'_a$ (eight (10×1)-vectors)

a_1

```
> M:=(y,z)->(t+y,z):
> aa[1]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_2

```
> M:=(y,z)->((1+t)*y,z):
> aa[2]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_3

```
> M:=(y,z)->(y+t*z,z):
> aa[3]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_4

```
> M:=(y,z)->(y,t+z):
> aa[4]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_5

```
> M:=(y,z)->(y,t*y+z):
> aa[5]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_6

```
> M:=(y,z)->(y,(1+t)*z):
> aa[6]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_7

```
> M:=(y,z)->(y/(1-t*y),z/(1-t*y)):
> aa[7]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

a_8

```
> M:=(y,z)->(y/(1-t*z),z/(1-t*z)):
> aa[8]:=MatrixMatrixMultiply(R_2,champ4(M)):
```

6.5.3 Columns of $\mathscr{M}''_a$ (eight (10×1)-vectors)

a_1

```
> M:=(y,z)->(t+y,z):
> aaa[1]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_2

```
> M:=(y,z)->((1+t)*y,z):
> aaa[2]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_3

```
> M:=(y,z)->(y+t*z,z):
> aaa[3]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_4

```
> M:=(y,z)->(y,t+z):
> aaa[4]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_5

```
> M:=(y,z)->(y,t*y+z):
> aaa[5]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_6

```
> M:=(y,z)->(y,(1+t)*z):
> aaa[6]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_7

```
> M:=(y,z)->(y/(1-t*y),z/(1-t*y)):
> aaa[7]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

a_8

```
> M:=(y,z)->(y/(1-t*z),z/(1-t*z)):
> aaa[8]:=MatrixMatrixMultiply(R_2,champ3(M)):
```

6.5.4 Construction of the matrix $\mathscr{V}_1$

```
>  mise:=proc(k::integer)
>  local u,v,w,h:
>  u:=convert(Transpose(a[k]),list):
>  v:=convert(Transpose(aa[k]),list):
>  w:=convert(Transpose(aaa[k]),list):
>  h:=[op(u),op(v),op(w)]:
>  RETURN(convert(h,Vector))
>  end:
>  for k from 1 to 8 do
>  VectDef[k]:=mise(k)
>  od:
```

6.6 The 63 vectors of $V(\mathfrak{D}_2)$ that form the columns of the matrix $\mathscr{V}_2$

6.6.1 Vectors of type a (we compute the columns of the matrices $\mathscr{N}_{\mathrm{a}}'$, $\mathscr{N}_{\mathrm{a}}$ et $\mathscr{N}_{\mathrm{a}}''$)

Columns of $\mathscr{N}_{\mathrm{a}}'$ *(eight* (10×1)*-vectors)*

Aa_1

```
>  M:=(y,z)->(t+y,z):
>  aA[1]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_2

```
>  M:=(y,z)->((1+t)*y,z):
>  aA[2]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_3

```
>  M:=(y,z)->(y+t*z,z):
>  aA[3]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_4

```
>  M:=(y,z)->(y,t+z):
>  aA[4]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_5

```
>  M:=(y,z)->(y,t*y+z):
>  aA[5]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_6

```
>  M:=(y,z)->(y,(1+t)*z):
>  aA[6]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_7

```
>  M:=(y,z)->(y/(1-t*y),z/(1-t*y)):
>  aA[7]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Aa_8

```
>  M:=(y,z)->(y/(1-t*z),z/(1-t*z)):
>  aA[8]:=MatrixMatrixMultiply(R_1,champ1(M)):
```

Columns of $\mathscr{N}_{\mathrm{a}}$ *to which we add 55 zeros*

Aa_1

```
>  pro1:=[-mu[1],-1,2,0,0,0,-mu[4]^2,-2*mu[4],0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro1);
>  aaA[1]:=convert(pro1,Vector):
```

$$pro1 := [-\mu_1, -1, 2, 0, 0, 0, -\mu_4^2, -2\,\mu_4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

$$65$$

Aa_2

```
>  pro2:=[0,0,0,0,0,0,3,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro2);
>  aaA[2]:=convert(pro2,Vector):
```

$$pro2 := [0, 0, 0, 0, 0, 0, 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0$$
$$, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_3

```
>  pro3:=[0,0,0,0,0,0,0,0,3,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro3);
>  aaA[3]:=convert(pro3,Vector):
```

$$pro3 := [0, 0, 0, 0, 0, 0, 0, 0, 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0$$
$$, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_4

```
>  pro4:=[1,0,0,0,0,0,-mu[4]^3,-3*mu[4]^2,2*mu[5]^2,4*mu[5],0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro4);
>  aaA[4]:=convert(pro4,Vector):
```

$$pro4 := [1, 0, 0, 0, 0, 0, -\mu_4^3, -3\,\mu_4^2, 2\,\mu_5^2, 4\,\mu_5, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$$
$$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_5

```
>  pro5:=[0,0,-mu[2],-1,-2*mu[3],-2,-3*mu[4],-3,-4*mu[5],-4,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro5);
>  aaA[5]:=convert(pro5,Vector):
```

$$pro5 := [0, 0, -\mu_2, -1, -2\,\mu_3, -2, -3\,\mu_4, -3, -4\,\mu_5, -4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$$
$$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_6

```
>  pro6:=[0,0,0,0,0,0,-2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro6);
>  aaA[6]:=convert(pro6,Vector):
```

$$pro6 := [0, 0, 0, 0, 0, 0, -2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0$$
$$, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_7

```
>  pro7:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro7);
>  aaA[7]:=convert(pro7,Vector):
```

$$pro7 := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$$
$$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Aa_8

```
>  pro8:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro8);
>  aaA[8]:=convert(pro7,Vector):
```

$$\mathit{pro8} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

Columns of $\mathscr{N}_{\mathrm{a}}''$ (*eight* (10×1)-*vectors*)

Aa_1

```
> M:=(y,z)->(t+y,z):
> aaaA[1]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_2

```
> M:=(y,z)->((1+t)*y,z):
> aaaA[2]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_3

```
> M:=(y,z)->(y+t*z,z):
> aaaA[3]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_4

```
> M:=(y,z)->(y,t+z):
> aaaA[4]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_5

```
> M:=(y,z)->(y,t*y+z):
> aaaA[5]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_6

```
> M:=(y,z)->(y,(1+t)*z):
> aaaA[6]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_7

```
> M:=(y,z)->(y/(1-t*y),z/(1-t*y)):
> aaaA[7]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Aa_8

```
> M:=(y,z)->(y/(1-t*z),z/(1-t*z)):
> aaaA[8]:=simplify(MatrixMatrixMultiply(R_2,champ2(M))):
```

Glueing

We form the first eight vectors of type a of $V(\mathfrak{D}_2)$.

```
> for k from 1 to 8 do
> for j from 1 to 10 do
> VectDef_a[k][j]:=aA[k][j]
> od
> od:
> for k from 1 to 8 do
> for j from 11 to 75 do
> VectDef_a[k][j]:=aaA[k][j-10]
> od
> od:
> for k from 1 to 8 do
> for j from 76 to 85 do
> VectDef_a[k][j]:=aaaA[k][j-75]
> od
> od:
> for k from 1 to 8 do
> VectDef_a[k]:=convert(VectDef_a[k],list)
> od:
> for k from 1 to 8 do
> VectDef_a[k]:=convert(VectDef_a[k],Vector)
> od:
```

6.6.2 Vectors of type b (we compute the columns of the matrices $\mathscr{N}_{\mathrm{b}}'$, $\mathscr{N}_{\mathrm{b}}$ and $\mathscr{N}_{\mathrm{b}}''$)

Columns of $\mathscr{N}_{\mathrm{b}}'$ *(twenty five* (10×1)*-vectors)*

```
> Mg:=(y,z,p,q)->(y+t*y^p/z^q,z);
```

$$Mg := (y, z, p, q) \mapsto y + \frac{t y^p}{z^q}, z$$

```
> M:=(y,z)->Mg(y,z,p,q);
```

$$M := (y, z) \mapsto Mg(y, z, p, q)$$

$Ab_{p,q}$

```
> bA:=unapply(MatrixMatrixMultiply(R_1,champ1(M)),(p,q)):
```

Colums of $\mathscr{N}_{\mathrm{b}}$ *to which we add a* (25×25)*-identity block then* 30 *zeros*

$Ab_{0,1}$

```
> pro1:=[0,0,0,0,0,0,2*mu[4]^5,10*mu[4]^4,3*mu[5]^3,9*mu[5]^2, 1,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro1);
> bbA[0,1]:=convert(pro1,Vector):
> bbA[0,1][11];
```

$pro1 := [0, 0, 0, 0, 0, 0, 2\,\mu_4^5, 10\,\mu_4^4, 3\,\mu_5^3, 9\,\mu_5^2, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{1,1}$

```
> pro2:=[-1,0,0,0,0,0,mu[4]^3,3*mu[4]^2,-2*mu[5]^2,-4*mu[5],0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro2);
> bbA[1,1]:=convert(pro2,Vector):
> bbA[1,1][12];
```

$pro2 := [-1, 0, 0, 0, 0, 0, \mu_4^3, 3\,\mu_4^2, -2\,\mu_5^2, -4\,\mu_5, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{2,1}$

```
> pro3:=[0,0,2*mu[2],2,3*mu[3],3,4*mu[4],4,5*mu[5],5, 0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro3);
> bbA[2,1]:=convert(pro3,Vector):
> bbA[2,1][13];
```

$pro3 := [0, 0, 2\,\mu_2, 2, 3\,\mu_3, 3, 4\,\mu_4, 4, 5\,\mu_5, 5, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{0,2}$

```
> pro4:=[0,0,0,0,0,0,-9*mu[4]^8,-72*mu[4]^7,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro4);
> bbA[0,2]:=convert(pro4,Vector):
> bbA[0,2][14];
```

$pro4 := [0, 0, 0, 0, 0, 0, -9\,\mu_4^8, -72\,\mu_4^7, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{1,2}$

```
>  pro5:=[0,0,0,0,0,0,-3*mu[4]^6,-18*mu[4]^5,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro5);
>  bbA[1,2]:=convert(pro5,Vector):
>  bbA[1,2][15];
```

$$\mathit{pro5} := [0, 0, 0, 0, 0, 0, -3\,\mu_4^6, -18\,\mu_4^5, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{2,2}$

```
>  pro6:=[-1/mu[1],1/mu[1]^2,0,0,0,0,-mu[4]^4,-4*mu[4]^3,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro6);
>  bbA[2,2]:=convert(pro6,Vector):
>  bbA[2,2][16];
```

$$\mathit{pro6} := [-\mu_1^{-1}, \mu_1^{-2}, 0, 0, 0, 0, -\mu_4^4, -4\,\mu_4^3, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{3,2}$

```
>  pro7:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro7);
>  bbA[3,2]:=convert(pro7,Vector):
>  bbA[3,2][17];
```

$$\mathit{pro7} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{0,3}$

```
>  pro8:=[0,0,0,0,0,0,52*mu[4]^11,572*mu[4]^10,-28*mu[5]^6,-168*mu[5]^5,0,0,0,0,0,0,0,1,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0];
>  nops(pro8);
>  bbA[0,3]:=convert(pro8,Vector):
>  bbA[0,3][18];
```

$$\mathit{pro8} := [0, 0, 0, 0, 0, 0, 52\,\mu_4^{11}, 572\,\mu_4^{10}, -28\,\mu_5^6, -168\,\mu_5^5, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{1,3}$

```
>  pro9:=[0,0,0,0,0,0,15*mu[4]^9,135*mu[4]^8,12*mu[5]^5,60*mu[5]^4,0,0,0,0,0,0,0,0,1,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro1);
>  bbA[1,3]:=convert(pro9,Vector):
>  bbA[1,3][19];
```

$$\mathit{pro9} := [0, 0, 0, 0, 0, 0, 15\,\mu_4^9, 135\,\mu_4^8, 12\,\mu_5^5, 60\,\mu_5^4, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{2,3}$

```
> pro10:=[0,0,0,0,0,0,4*mu[4]^7,28*mu[4]^6,-5*mu[5]^4,-20*mu[5]^3,0,0,0,0,0,0,0,0,0,1,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro10);
> bbA[2,3]:=convert(pro10,Vector):
> bbA[2,3][20];
```

$\mathit{pro10} := [0, 0, 0, 0, 0, 0, 4\,\mu_4^7, 28\,\mu_4^6, -5\,\mu_5^4, -20\,\mu_5^3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{3,3}$

```
> pro11:=[-1/mu[1]^2,2/mu[1]^3,0,0,0,0,mu[4]^5,5*mu[4]^4,2*mu[5]^3,6*mu[5]^2,0,0,0,0,0,0,
0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0];
> nops(pro11);
> bbA[3,3]:=convert(pro11,Vector):
> bbA[3,3][21];
```

$\mathit{pro11} := [-\mu_1^{-2}, 2\,\mu_1^{-3}, 0, 0, 0, 0, \mu_4^5, 5\,\mu_4^4, 2\,\mu_5^3, 6\,\mu_5^2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{4,3}$

```
> pro12:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro12);
> bbA[4,3]:=convert(pro12,Vector):
> bbA[4,3][22];
```

$\mathit{pro12} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
0
$, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{0,4}$

```
> pro13:=[0,0,0,0,0,0,-340*mu[4]^14,-4760*mu[4]^13,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro13);
> bbA[0,4]:=convert(pro13,Vector):
> bbA[0,4][23];
```

$\mathit{pro13} := [0, 0, 0, 0, 0, 0, -340\,\mu_4^{14}, -4760\,\mu_4^{13}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{1,4}$

```
> pro14:=[0,0,0,0,0,0,-91*mu[4]^12,-1092*mu[4]^11,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro14);
> bbA[1,4]:=convert(pro14,Vector):
> bbA[1,4][24];
```

$\mathit{pro14} := [0, 0, 0, 0, 0, 0, -91\,\mu_4^{12}, -1092\,\mu_4^{11}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$

65

1

$Ab_{2,4}$

```
> pro15:=[0,0,0,0,0,0,-22*mu[4]^10,-220*mu[4]^9,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro15);
> bbA[2,4]:=convert(pro15,Vector):
> bbA[2,4][25];
```

$$\mathit{pro15} := [0,0,0,0,0,0,-22\,\mu_4^{10},-220\,\mu_4^{9},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ab_{3,4}$

```
> pro16:=[0,0,0,0,0,0,-5*mu[4]^8,-40*mu[4]^7,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro16);
> bbA[3,4]:=convert(pro16,Vector):
> bbA[3,4][26];
```

$$\mathit{pro16} := [0,0,0,0,0,0,-5\,\mu_4^{8},-40\,\mu_4^{7},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ab_{4,4}$

```
> pro17:=[-1/mu[1]^3,3/mu[1]^4,0,0,0,0,-mu[4]^6,-6*mu[4]^5,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro17);
> bbA[4,4]:=convert(pro17,Vector):
> bbA[4,4][27];
```

$$\mathit{pro17} := [-\mu_1^{-3},3\,\mu_1^{-4},0,0,0,0,-\mu_4^{6},-6\,\mu_4^{5},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ab_{5,4}$

```
> pro18:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro18);
> bbA[5,4]:=convert(pro18,Vector):
> bbA[5,4][28];
```

$$\mathit{pro18} := [0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ab_{0,5}$

```
> pro19:=[0,0,0,0,0,0,2394*mu[4]^17,40698*mu[4]^16,429*mu[5]^9,3861*mu[5]^8,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0];
> nops(pro19);
> bbA[0,5]:=convert(pro19,Vector):
> bbA[0,5][29];
```

$$\mathit{pro19} := [0,0,0,0,0,0,2394\,\mu_4^{17},40698\,\mu_4^{16},429\,\mu_5^{9},3861\,\mu_5^{8},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ab_{1,5}$

```
>  pro20:=[0,0,0,0,0,0,612*mu[4]^15,9180*mu[4]^14,-165*mu[5]^8,-1320*mu[5]^7,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0];
>  nops(pro20);
>  bbA[1,5]:=convert(pro20,Vector):
>  bbA[1,5][30];
```

$$pro20\ :=\ [0, 0, 0, 0, 0, 0, 612\,\mu_4^{15}, 9180\,\mu_4^{14}, -165\,\mu_5^{8}, -1320\,\mu_5^{7}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{2,5}$

```
>  pro21:=[0,0,0,0,0,0,140*mu[4]^13,1820*mu[4]^12,60*mu[5]^7,420*mu[5]^6,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0];
>  nops(pro21);
>  bbA[2,5]:=convert(pro21,Vector):
>  bbA[2,5][31];
```

$$pro21\ :=\ [0, 0, 0, 0, 0, 0, 140\,\mu_4^{13}, 1820\,\mu_4^{12}, 60\,\mu_5^{7}, 420\,\mu_5^{6}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{3,5}$

```
>  pro22:=[0,0,0,0,0,0,30*mu[4]^11,330*mu[4]^10,-21*mu[5]^6,-126*mu[5]^5,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0];
>  nops(pro22);
>  bbA[3,5]:=convert(pro22,Vector):
>  bbA[3,5][32];
```

$$pro22\ :=\ [0, 0, 0, 0, 0, 0, 30\,\mu_4^{11}, 330\,\mu_4^{10}, -21\,\mu_5^{6}, -126\,\mu_5^{5}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{4,5}$

```
>  pro23:=[0,0,0,0,0,0,6*mu[4]^9,54*mu[4]^8,7*mu[5]^5,35*mu[5]^4,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro23);
>  bbA[4,5]:=convert(pro23,Vector):
>  bbA[4,5][33];
```

$$pro23\ :=\ [0, 0, 0, 0, 0, 0, 6\,\mu_4^{9}, 54\,\mu_4^{8}, 7\,\mu_5^{5}, 35\,\mu_5^{4}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{5,5}$

```
>  pro24:=[-1/mu[1]^4,4/mu[1]^5,0,0,0,0,mu[4]^7,7*mu[4]^6,-2*mu[5]^4,-8*mu[5]^3,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0];
>  nops(pro24);
>  bbA[5,5]:=convert(pro24,Vector):
>  bbA[5,5][34];
```

$$pro24\ :=\ [-\mu_1^{-4}, 4\,\mu_1^{-5}, 0, 0, 0, 0, \mu_4^{7}, 7\,\mu_4^{6}, -2\,\mu_5^{4}, -8\,\mu_5^{3}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ab_{6,5}$

```
>  pro25:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro25);
>  bbA[6,5]:=convert(pro25,Vector):
>  bbA[6,5][35];
```

$pro25 := [0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,$
$0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$

65

1

Columns of $\mathscr{N}_{\mathrm{b}}''$ (*twenty five* (10×1)*-vectors*)

```
>  Mg:=(y,z,p,q)->(y+t*y^p/z^q,z);
```

$$Mg := (y,z,p,q) \mapsto y+\frac{ty^p}{z^q},z$$

```
>  M:=(y,z)->Mg(y,z,p,q);
```

$$M := (y,z) \mapsto Mg(y,z,p,q)$$

$Ab_{p,q}$

```
>  bbbA:=unapply(MatrixMatrixMultiply(R_2,champ2(M)),(p,q)):
```

Glueing
We form the 25 vectors of type b of $V(\mathfrak{D}_2)$.

```
>  for q from 1 to 5 do
>  for p from 0 to q+1 do
>  for j from 1 to 10 do
>  u[p,q][j]:=bA(p,q)[j]
>  od
>  od
>  od:
>  for q from 1 to 5 do
>  for p from 0 to q+1 do
>  for j from 11 to 75 do
>  u[p,q][j]:=bbA[p,q][j-10]
>  od
>  od
>  od:
>  for q from 1 to 5 do
>  for p from 0 to q+1 do
>  for j from 76 to 85 do
>  u[p,q][j]:=bbbA(p,q)[j-75]
>  od
>  od
>  od:
>  for q from 1 to 5 do
>  for p from 0 to q+1 do
>  u[p,q]:=convert(u[p,q],list)
>  od
>  od:
>  q:=1:
>  for p from 0 to q+1 do
>  VectDef_b[p+1]:=u[p,q]
>  od:
>  q:=2:
>  for p from 0 to q+1 do
>  VectDef_b[p+4]:=u[p,q]
>  od:
>  q:=3:
>  for p from 0 to q+1 do
>  VectDef_b[p+8]:=u[p,q]
>  od:
>  q:=4:
>  for p from 0 to q+1 do
>  VectDef_b[p+13]:=u[p,q]
>  od:
>  q:=5:
>  for p from 0 to q+1 do
>  VectDef_b[p+19]:=u[p,q]
>  od:
>  p:='p':
>  q:='q':
```

6.6.3 Vectors of type c (we compute the columns of the matrices $\mathscr{N}_{\mathrm{c}}'$, $\mathscr{N}_{\mathrm{c}}$ and $\mathscr{N}_{\mathrm{c}}''$)

Columns of $\mathscr{N}_{\mathrm{c}}'$ *(twenty five* (10×1)*-vectors)*

```
>  Mg:=(y,z,p,q)->(y,z+t*y^p/z^q);
```

$$Mg := (y, z, p, q) \mapsto y, z + \frac{t y^p}{z^q}$$

```
>  M:=(y,z)->Mg(y,z,p,q);
```

$$M := (y, z) \mapsto Mg(y, z, p, q)$$

$Ac_{p,q}$

```
>  cA:=unapply(MatrixMatrixMultiply(R_1,champ1(M)),(p,q)):
```

Columns of $\mathscr{N}_{\mathrm{c}}$ *to which we add* 25 *zeros, a* (25×25)*-identity block then* 5 *zeros*

$Ac_{0,1}$

```
>  pro1:=[0,0,0,0,0,0,4*mu[4]^6,24*mu[4]^5,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro1);
>  ccA[0,1]:=convert(pro1,Vector):
>  ccA[0,1][36];
```

$$pro1 := [0,0,0,0,0,0,4\,\mu_4^6,24\,\mu_4^5,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{1,1}$

```
>  pro2:=[1/mu[1],-1/mu[1]^2,0,0,0,0,mu[4]^4,4*mu[4]^3,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro2);
>  ccA[1,1]:=convert(pro2,Vector):
>  ccA[1,1][37];
```

$$pro2 := [\mu_1^{-1},-\mu_1^{-2},0,0,0,0,\mu_4^4,4\,\mu_4^3,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{2,1}$

```
>  pro3:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro3);
>  ccA[2,1]:=convert(pro3,Vector):
>  ccA[2,1][38];
```

$$pro3 := [0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{0,2}$

```
>  pro4:=[0,0,0,0,0,0,-25*mu[4]^9,-225*mu[4]^8,-18*mu[5]^5,-90*mu[5]^4,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
```

```
>  nops(pro4);
>  ccA[0,2]:=convert(pro4,Vector):
>  ccA[0,2][39];
```

$$\mathit{pro4} := [0, 0, 0, 0, 0, 0, -25\,\mu_4^9, -225\,\mu_4^8, -18\,\mu_5^5, -90\,\mu_5^4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{1,2}$

```
>  pro5:=[0,0,0,0,0,0,-5*mu[4]^7,-35*mu[4]^6,6*mu[5]^4,24*mu[5]^3,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro5);
>  ccA[1,2]:=convert(pro5,Vector):
>  ccA[1,2][40];
```

$$\mathit{pro5} := [0, 0, 0, 0, 0, 0, -5\,\mu_4^7, -35\,\mu_4^6, 6\,\mu_5^4, 24\,\mu_5^3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{2,2}$

```
>  pro6:=[1/mu[1]^2,-2/mu[1]^3,0,0,0,0,-mu[4]^5,-5*mu[4]^4,-2*mu[5]^3,-6*mu[5]^2,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0];
>  nops(pro6);
>  ccA[2,2]:=convert(pro6,Vector):
>  ccA[2,2][41];
```

$$\mathit{pro6} := [\mu_1^{-2}, -2\,\mu_1^{-3}, 0, 0, 0, 0, -\mu_4^5, -5\,\mu_4^4, -2\,\mu_5^3, -6\,\mu_5^2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{3,2}$

```
>  pro7:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro7);
>  ccA[3,2]:=convert(pro7,Vector):
>  ccA[3,2][42];
```

$$\mathit{pro7} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{0,3}$

```
>  pro8:=[0,0,0,0,0,0,182*mu[4]^12,2184*mu[4]^11,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro8);
>  ccA[0,3]:=convert(pro8,Vector):
>  ccA[0,3][43];
```

$$\mathit{pro8} := [0, 0, 0, 0, 0, 0, 182\,\mu_4^{12}, 2184\,\mu_4^{11}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{1,3}$

```
>  pro9:=[0,0,0,0,0,0,33*mu[4]^10,330*mu[4]^9,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
```

```
>  nops(pro1);
>  ccA[1,3]:=convert(pro9,Vector):
>  ccA[1,3][44];
```

$$\mathit{pro9} := [0,0,0,0,0,0,33\,\mu_4^{10},330\,\mu_4^{9},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{2,3}$

```
>  pro10:=[0,0,0,0,0,0,6*mu[4]^8,48*mu[4]^7,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro10);
>  ccA[2,3]:=convert(pro10,Vector):
>  ccA[2,3][45];
```

$$\mathit{pro10} := [0,0,0,0,0,0,6\,\mu_4^{8},48\,\mu_4^{7},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{3,3}$

```
>  pro11:=[1/mu[1]^3,-3/mu[1]^4,0,0,0,0,mu[4]^6,6*mu[4]^5,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro11);
>  ccA[3,3]:=convert(pro11,Vector):
>  ccA[3,3][46];
```

$$\mathit{pro11} := [\mu_1^{-3},-3\,\mu_1^{-4},0,0,0,0,\mu_4^{6},6\,\mu_4^{5},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{4,3}$

```
>  pro12:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro12);
>  ccA[4,3]:=convert(pro12,Vector):
>  ccA[4,3][47];
```

$$\mathit{pro12} := [0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{0,4}$

```
>  pro13:=[0,0,0,0,0,0,-1428*mu[4]^15,-21420*mu[4]^14,330*mu[5]^8,2640*mu[5]^7,
>  0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro13);
>  ccA[0,4]:=convert(pro13,Vector):
>  ccA[0,4][48];
```

$$\mathit{pro13} := [0,0,0,0,0,0,-1428\,\mu_4^{15},-21420\,\mu_4^{14},330\,\mu_5^{8},2640\,\mu_5^{7},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0]$$

65

1

$Ac_{1,4}$

```
>  pro14:=[0,0,0,0,0,0,-245*mu[4]^13,-3185*mu[4]^12,-96*mu[5]^7,-672*mu[5]^6,
>  0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0];
```

```
>  nops(pro14);
>  ccA[1,4]:=convert(pro14,Vector):
>  ccA[1,4][49];
```

$$\mathit{pro14} := [0, 0, 0, 0, 0, 0, -245\,\mu_4^{13}, -3185\,\mu_4^{12}, -96\,\mu_5^{7}, -672\,\mu_5^{6}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{2,4}$

```
>  pro15:=[0,0,0,0,0,0,-42*mu[4]^11,-462*mu[4]^10,28*mu[5]^6,168*mu[5]^5,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0];
>  nops(pro15);
>  ccA[2,4]:=convert(pro15,Vector):
>  ccA[2,4][50];
```

$$\mathit{pro15} := [0, 0, 0, 0, 0, 0, -42\,\mu_4^{11}, -462\,\mu_4^{10}, 28\,\mu_5^{6}, 168\,\mu_5^{5}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{3,4}$

```
>  pro16:=[0,0,0,0,0,0,-7*mu[4]^9,-63*mu[4]^8,-8*mu[5]^5,-40*mu[5]^4,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro16);
>  ccA[3,4]:=convert(pro16,Vector):
>  ccA[3,4][51];
```

$$\mathit{pro16} := [0, 0, 0, 0, 0, 0, -7\,\mu_4^{9}, -63\,\mu_4^{8}, -8\,\mu_5^{5}, -40\,\mu_5^{4}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{4,4}$

```
>  pro17:=[1/mu[1]^4,-4/mu[1]^5,0,0,0,0,-mu[4]^7,-7*mu[4]^6,2*mu[5]^4,8*mu[5]^3,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,
0,0,0,0,0];
>  nops(pro17);
>  ccA[4,4]:=convert(pro17,Vector):
>  ccA[4,4][52];
```

$$\mathit{pro17} := [\mu_1^{-4}, -4\,\mu_1^{-5}, 0, 0, 0, 0, -\mu_4^{7}, -7\,\mu_4^{6}, 2\,\mu_5^{4}, 8\,\mu_5^{3}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{5,4}$

```
>  pro18:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0];
>  nops(pro18);
>  ccA[5,4]:=convert(pro18,Vector):
>  ccA[5,4][53];
```

$$\mathit{pro18} := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$$

65

1

$Ac_{0,5}$

```
> pro19:=[0,0,0,0,0,0,11704*mu[4]^18,210672*mu[4]^17,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0];
> nops(pro19);
> ccA[0,5]:=convert(pro19,Vector):
> ccA[0,5][54];
```

$$pro19 := [0,0,0,0,0,0,11704\,\mu_4^{18},210672\,\mu_4^{17},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0]$$

$$65$$

$$1$$

$$Ac_{1,5}$$

```
> pro20:=[0,0,0,0,0,0,1938*mu[4]^16,31008*mu[4]^15,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0];
> nops(pro20);
> ccA[1,5]:=convert(pro20,Vector):
> ccA[1,5][55];
```

$$pro20 := [0,0,0,0,0,0,1938\,\mu_4^{16},31008\,\mu_4^{15},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0]$$

$$65$$

$$1$$

$$Ac_{2,5}$$

```
> pro21:=[0,0,0,0,0,0,320*mu[4]^14,4480*mu[4]^13,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0];
> nops(pro21);
> ccA[2,5]:=convert(pro21,Vector):
> ccA[2,5][56];
```

$$pro21 := [0,0,0,0,0,0,320\,\mu_4^{14},4480\,\mu_4^{13},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0]$$

$$65$$

$$1$$

$$Ac_{3,5}$$

```
> pro22:=[0,0,0,0,0,0,52*mu[4]^12,624*mu[4]^11,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0];
> nops(pro22);
> ccA[3,5]:=convert(pro22,Vector):
> ccA[3,5][57];
```

$$pro22 := [0,0,0,0,0,0,52\,\mu_4^{12},624\,\mu_4^{11},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0]$$

$$65$$

$$1$$

$$Ac_{4,5}$$

```
> pro23:=[0,0,0,0,0,0,8*mu[4]^10,80*mu[4]^9,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0];
> nops(pro23);
> ccA[4,5]:=convert(pro23,Vector):
> ccA[4,5][58];
```

$$pro23 := [0,0,0,0,0,0,8\,\mu_4^{10},80\,\mu_4^{9},0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0]$$

$$65$$

$$1$$

$$Ac_{5,5}$$

```
> pro24:=[1/mu[1]^5,-5/mu[1]^6,0,0,0,0,mu[4]^8,8*mu[4]^7,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0];
> nops(pro24);
> ccA[5,5]:=convert(pro24,Vector):
> ccA[5,5][59];
```

$$pro24 := [\mu_1^{-5}, -5\,\mu_1^{-6}, 0, 0, 0, 0, \mu_4^{8}, 8\,\mu_4^{7}, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0]$$

65

1

Ac$_{6,5}$

```
> pro25:=[0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0];
> nops(pro25);
> ccA[6,5]:=convert(pro25,Vector):
> ccA[6,5][60];
```

$$pro25 := [0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0]$$

65

1

Columns of $\mathscr{N}_c''$ (*twenty five* (10 × 1)*-vectors*)

```
> Mg:=(y,z,p,q)->(y,z+t*y^p/z^q);
```

$$Mg := (y, z, p, q) \mapsto y, z + \frac{t y^p}{z^q}$$

```
> M:=(y,z)->Mg(y,z,p,q);
```

$$M := (y, z) \mapsto Mg(y, z, p, q)$$

Ac$_{p,q}$

```
> cccA:=unapply(MatrixMatrixMultiply(R_2,champ2(M)),(p,q)):
```

Glueing
We form the 25 vectors of type c of $V(\mathfrak{D}_2)$.

```
> for q from 1 to 5 do
> for p from 0 to q+1 do
> for j from 1 to 10 do
> uu[p,q][j]:=cA(p,q)[j]
> od
> od
> od:
> for q from 1 to 5 do
> for p from 0 to q+1 do
> for j from 11 to 75 do
> uu[p,q][j]:=ccA[p,q][j-10]
> od
> od
> od:
> for q from 1 to 5 do
> for p from 0 to q+1 do
> for j from 76 to 85 do
> uu[p,q][j]:=cccA(p,q)[j-75]
> od
> od
> od:
> for q from 1 to 5 do
> for p from 0 to q+1 do
> uu[p,q]:=convert(uu[p,q],list)
> od
> od:
> q:=1:
> for p from 0 to q+1 do
> VectDef_c[p+1]:=uu[p,q]
> od:
```

```
>  q:=2:
>  for p from 0 to q+1 do
>  VectDef_c[p+4]:=uu[p,q]
>  od:
>  q:=3:
>  for p from 0 to q+1 do
>  VectDef_c[p+8]:=uu[p,q]
>  od:
>  q:=4:
>  for p from 0 to q+1 do
>  VectDef_c[p+13]:=uu[p,q]
>  od:
>  q:=5:
>  for p from 0 to q+1 do
>  VectDef_c[p+19]:=uu[p,q]
>  od:
>  p:='p':
>  q:='q':
```

6.6.4 Vectors of type d (we compute the columns of the matrices $\mathscr{N}_{\rm d}'$, $\mathscr{N}_{\rm d}$ and $\mathscr{N}_{\rm d}''$)

Columns of $\mathscr{N}_{\rm d}'$ *(five* (10×1)*-vectors)*

```
>  Mg:=(y,z,p)->(y+t*y^(p+2)/z^p,z+t*y^(p+1)/z^(p-1));
```

$$Mg := (y, z, p) \mapsto y + \frac{t y^{p+2}}{z^p}, z + \frac{t y^{p+1}}{z^{p-1}}$$

```
>  M:=(y,z)->Mg(y,z,p);
```

$$M := (y, z) \mapsto Mg(y, z, p)$$

Ad_p

```
>  dA:=unapply(MatrixMatrixMultiply(R_1,champ1(M)),p):
```

Columns of $\mathscr{N}_{\rm d}$ *to which we add* 50 *zeros then a* (5×5)*-identity block*

Ad_1

```
>  pro1:=[0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0];
>  nops(pro1);
>  ddA[1]:=convert(pro1,Vector):
>  ddA[1][61];
```

$$pro1 := [0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$$
$$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, 0]$$

$$65$$

$$1$$

Ad_2

```
>  pro2:=[0,0,mu[2]^2,2*mu[2],0,0,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0];
>  nops(pro2);
>  ddA[2]:=convert(pro2,Vector):
>  ddA[2][62];
```

$$pro2 := [0, 0, \mu_2^2, 2\,\mu_2, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$$
$$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0]$$

$$65$$

$$1$$

Ad_3

```
>  pro3:=[0,0,0,0,mu[3]^2,2*mu[3],2*mu[4],2,2*mu[5],2,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0];
>  nops(pro3);
>  ddA[3]:=convert(pro3,Vector):
>  ddA[3][63];
```

$pro3 := [0, 0, 0, 0, \mu_3^2, 2\,\mu_3, 2\,\mu_4, 2, 2\,\mu_5, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0]$

65

1

Ad$_4$

```
>  pro4:=[0,0,0,0,0,0,mu[4]^2,2*mu[4],0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0];
>  nops(pro4);
>  ddA[4]:=convert(pro4,Vector):
>  ddA[4][64];
```

$pro4 := [0, 0, 0, 0, 0, 0, \mu_4^2, 2\,\mu_4, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0]$

65

1

Ad$_5$

```
>  pro5:=[0,0,0,0,0,0,0,0,mu[5]^2,2*mu[5],0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,
0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1];
>  nops(pro5);
>  ddA[5]:=convert(pro5,Vector):
>  ddA[5][65];
```

$pro5 := [0, 0, 0, 0, 0, 0, 0, 0, \mu_5^2, 2\,\mu_5, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,$
$0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1]$

65

1

Columns of $\mathcal{N}_{\mathrm{d}}''$ *(five* (10×1)*-vectors)*

```
>  Mg:=(y,z,p)->(y+t*y^(p+2)/z^p,z+t*y^(p+1)/z^(p-1));
```

$$Mg := (y, z, p) \mapsto y + \frac{t y^{p+2}}{z^p}, z + \frac{t y^{p+1}}{z^{p-1}}$$

```
>  M:=(y,z)->Mg(y,z,p);
```

$$M := (y, z) \mapsto Mg(y, z, p)$$

Ad$_p$

```
>  dddA:=unapply(MatrixMatrixMultiply(R_2,champ2(M)),p):
```

Glueing
We form the 5 vectors of type d of $V(\mathfrak{D}_2)$.

```
>  for k from 1 to 5 do
>  for j from 1 to 10 do
>  VectDef_d[k][j]:=dA(k)[j]
>  od
>  od:
>  for k from 1 to 5 do
>  for j from 11 to 75 do
>  VectDef_d[k][j]:=ddA[k][j-10]
>  od
>  od:
>  for k from 1 to 5 do
>  for j from 76 to 85 do
>  VectDef_d[k][j]:=dddA(k)[j-75]
>  od
>  od:
>  for k from 1 to 5 do
>  VectDef_d[k]:=convert(VectDef_d[k],list)
>  od:
>  for k from 1 to 5 do
>  VectDef_d[k]:=convert(VectDef_d[k],Vector)
>  od:
```

6.6.5 Construction of the matrix $\mathscr{V}_2$

We put together the 8 vectors of type a, the 25 vectors of type b, the 25 vectors of type c and the 5 vectors of type d.

```
>  for k from 1 to 8 do
>  V[k]:=VectDef_a[k]
>  od:
>  for k from 9 to 33 do
>  V[k]:=VectDef_b[k-8]
>  od:
>  for k from 34 to 58 do
>  V[k]:=VectDef_c[k-33]
>  od:
>  for k from 59 to 63 do
>  V[k]:=VectDef_d[k-58]
>  od:
>  for k from 1 to 63 do
>  WW[k]:=convert(V[k],list)
>  od:
>  MatVectDef:=Transpose(Matrix([seq(WW[k],k=1..63)])):
```

We define the images of the 8 vectors of $V(\mathfrak{D}_1)$ under the action of $\mathscr{Y}$.

```
>  for k from 1 to 8 do
>  ImageVectDef[k]:=MatrixMatrixMultiply(MM,VectDef[k])
>  od:
```

We know how to compute these 8 vectors in the basis with the 65 vectors of $V(\mathfrak{D}_2)$ thanks to the relations (5.3) given in the article. We check that these relations are satisfied (these 8 instructions give 85 zeros).

```
>  for k from 1 to 85 do
>  simplify(ImageVectDef[1][k]-WW[1][k]-3*WW[62][k]+4*WW[15][k]+3*WW[36][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[2][k]-WW[2][k]+3*WW[60][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[3][k]-WW[3][k]+3*WW[59][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[4][k]-WW[4][k]-2*WW[20][k]-WW[40][k]+2*WW[63][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[5][k]-WW[5][k]-2*WW[61][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[6][k]-WW[6][k]-2*WW[60][k]);
>  od:
>  for k from 1 to 85 do
>  simplify(ImageVectDef[7][k]-WW[7][k]);
>  od:
```

6.7 Computation of the action of ψ_* on $\mathrm{H}^1(X, \mathrm{T}X)$

Construction of the supplementary $\mathscr{E}_1$ of $V(\mathfrak{D}_1)$ of dimension 22 in $W(\mathfrak{D}_1)$.

We form a (30×30)-matrix $\mathtt{NN}_1$ whose the first 22 columns are now zero and the last 8 are the basis vectors of $V(\mathfrak{D}_1)$ computed previously.

```
>  for k from 1 to 30 do
>  for j from 23 to 30 do
>  N1[k][j]:=VectDef[j-22][k]
>  od
>  od:
>  for k from 1 to 30 do
>  for j from 1 to 22 do
>  N1[k][j]:=0
>  od
>  od:
>  for k from 1 to 30 do
>  N1[k]:=convert(N1[k],list)
>  od:
>  NN1:=Matrix([seq(N1[k],k=1..30)]):
```

We fill the first 22 columns of $\mathtt{NN}_1$ by some vectors chosen among the algebraic basis of $W(\mathfrak{D}_1)$; they will define a supplementary $\mathscr{E}_1$ of $V(\mathfrak{D}_1)$.

```
>  NN1[1,1]:=1:
>  NN1[2,2]:=1:
>  NN1[3,3]:=1:
>  NN1[4,4]:=1:
>  NN1[5,5]:=1:
>  NN1[6,6]:=1:
>  NN1[7,7]:=1:
>  NN1[8,8]:=1:
>  NN1[9,9]:=1:
>  NN1[10,10]:=1:
>  NN1[11,11]:=1:
>  NN1[12,12]:=1:
>  NN1[13,13]:=1:
>  NN1[14,14]:=1:
>  NN1[15,15]:=1:
>  NN1[16,16]:=1:
>  NN1[17,17]:=1:
>  NN1[21,18]:=1:
>  NN1[22,19]:=1:
>  NN1[23,20]:=1:
>  NN1[24,21]:=1:
>  NN1[25,22]:=1:
```

We compute the determinant of $\mathtt{NN}_1$, that is nonzero for generic parameters μ_i; this shows that $\mathscr{E}_1$ is a supplementary to $V(\mathfrak{D}_1)$.

```
>  Determinant(NN1);
```

$$12\,\mu_4\mu_5(-6\,\mu_4\mu_5 - 4\,\mu_4\mu_5^2 + 9\,\mu_5\alpha + 9\,\mu_4^2\alpha + 45\,\mu_5 + 45\,\mu_4^2 - 36\,\mu_4)$$

We extract the first 22 columns of $\mathtt{NN}_1$ in order to form the matrix `left`, that is the injection matrix of $\mathscr{E}_1$ into $W(\mathfrak{D}_1)$.

```
>  for k from 1 to 30 do
>  for j from 1 to 22 do
>  coeff1[k][j]:=NN1[k,j]:
>  od:
>  od:
>  for k from 1 to 30 do
>  coeff1[k]:=convert(coeff1[k],list)
>  od:
>  left:=Matrix([seq(coeff1[k],k=1..30)]):
```

Definition of the supplementary $\mathscr{E}_2$ of $V(\mathfrak{D}_2)$ in $W(\mathfrak{D}_2)$ of dimension 22 which is the image of $\mathscr{E}_1$ by the inclusion of $W(\mathfrak{D}_1)$ into $W(\mathfrak{D}_2)$.

We form a (85×85)-matrix $\mathtt{NN}_2$ whose the first 22 columns are now zero and the last 63 are some basis vectors of $V(\mathfrak{D}_2)$ computed previously.

```
>  for k from 1 to 85 do
>  for j from 23 to 85 do
>  N2[k][j]:=WW[j-22][k]
>  od
>  od:
>  for k from 1 to 85 do
>  for j from 1 to 22 do
>  N2[k][j]:=0
>  od
>  od:
>  for k from 1 to 85 do
>  N2[k]:=convert(N2[k],list)
>  od:
>  NN2:=Matrix([seq(N2[k],k=1..85)]):
```

We fill the first 22 columns of $\mathtt{NN}_2$ by the basis vectors of $\mathscr{E}_2$.

```
>  NN2[1,1]:=1:
>  NN2[2,2]:=1:
>  NN2[3,3]:=1:
>  NN2[4,4]:=1:
>  NN2[5,5]:=1:
```

```
> NN2[6,6]:=1:
> NN2[7,7]:=1:
> NN2[8,8]:=1:
> NN2[9,9]:=1:
> NN2[10,10]:=1:
> NN2[11,11]:=1:
> NN2[12,12]:=1:
> NN2[13,13]:=1:
> NN2[14,14]:=1:
> NN2[15,15]:=1:
> NN2[16,16]:=1:
> NN2[17,17]:=1:
> NN2[76,18]:=1:
> NN2[77,19]:=1:
> NN2[78,20]:=1:
> NN2[79,21]:=1:
> NN2[80,22]:=1:
```

The matrix `right` is the projection matrix of $W(\mathfrak{D}_2)$ onto $\mathscr{E}_2$.

```
> NNN2:=MatrixInverse(NN2):
> for k from 1 to 22 do
> for j from 1 to 85 do
> coeff2[k][j]:=NNN2[k,j]:
> od:
> od:
> for k from 1 to 22 do
> coeff2[k]:=convert(coeff2[k],list)
> od:
> right:=Matrix([seq(coeff2[k],k=1..22)]):
```

The matrix `OMG` is the matrix of f_* acting on $\mathrm{H}^1(X, \mathrm{T}X)$.

```
> OMG:=MatrixMatrixMultiply(right,MatrixMatrixMultiply(MM,left)):
> OMG;
```

```
[   ‘ 22 x 22 ‘ (Matrix)     ]
[                            ]
[ ‘Data Type:   ‘ anything   ]
[                            ]
[ ‘Storage:   ‘ rectangular  ]
[                            ]
[ ‘Order:   ‘ Fortran_order  ]
```

We assign the parameters λ_i et μ_i otherwise the computation is too heavy.

```
> lambda[1]:=10:
> lambda[2]:=-6:
> lambda[3]:=5:
> lambda[4]:=2:
> lambda[5]:=3:
> mu[1]:=3:
> mu[2]:=-2:
> mu[3]:=4:
> mu[4]:=3:
> mu[5]:=29:
```

Computation of the eigenvalues of `OMG`.

```
> factor(CharacteristicPolynomial(OMG,x));
```

$$(x^2+3\,x+1)(x^2+18\,x+1)(x^2-7\,x+1)(x^2+x+1)(x-1)^2(x+1)^4(x^2-x+1)^4$$

We look at the size of the Jordan blocks for the multiple eigenvalues.

```
> Id:=Matrix(22,22,shape=identity):
```

```
> Rank(OMG+Id);
> Rank((OMG+Id)^2);
> Rank((OMG+Id)^3);
```

```
19

18

18
```

```
>  Rank(OMG-Id);
>  Rank((OMG-Id)^2);

20

20

>  Rank(OMG-(1+I*sqrt(3))/2*Id);
>  Rank((OMG-(1+I*sqrt(3))/2*Id^2));

19

19
```

It takes 13 *minutes to execute the file.*

JULIE DÉSERTI, Universität Basel, Mathematisches Institut, Rheinsprung 21, CH-4051 Basel, Switzerland.
E-mail : `julie.deserti@unibas.ch` • On leave from Institut de Mathématiques de Jussieu, UMR 7586, Université Paris 7, Site Chevaleret, Case 7012, 75205 Paris Cedex 13, France. • *E-mail :* `deserti@math.jussieu.fr`

JULIEN GRIVAUX, CNRS, LATP, UMR 7353, CMI, Université de Provence, 39, rue Frédéric Joliot-Curie, 13453 Marseille Cedex 13, France.
E-mail : `jgrivaux@cmi.univ-mrs.fr`